NEW SOUTH WALES.

VITAL STATISTICS.

TWENTY-FIRST ANNUAL REPORT


FROM

THE REGISTRAR GENERAL, ON VITAL STATISTICS.

Presented to Parliament by Command.



SYDNEY: CHARLES POTTER, ACTING GOVERNMENT PRINTER.



VITAL STATISTICS.

THE REGISTRAR GENERAL TO THE HONORABLE THE COLONIAL SECRETARY.

TRANSMITTING ABSTRACTS OF

MARRIAGES, BIRTHS, AND DEATHS,

FOR THE YEAR 1876.

WITH PREFATORY REPORT.

Registrar General's Office, Sydney, 18 April, 1877.

SIR,

I have the honor to forward herewith the Returns of Marriages, Births, and Deaths, for the Population year ended 31st December, 1876. At this date the estimated population of the Colony numbered 629,776 souls, comprising 347,869 males, and 281,907 females. The total increase in population at the end of the year is estimated at 23,124 persons, or 3.81 per cent. The increase for the first six months of the year amounted to 10,514 persons and for the remaining period of the year to 12,610 persons. Of the total population the number of males was 65,962 in excess of females; the percentage for the former being 55.24, and for the latter 44.76, a difference of 10.48 in favour of males. To every 1,000 persons then there were 552 males and 448 females, which shows an increase of 2 males in the 1,000 on the previous year. For the first six months of the year there were 11,335 births, and 17,588 persons who arrived in the Colony by sea, a gross total of 28,923, from which I deduct 6,170 deaths for the same period, and the departures by sea 12,239, in all 18,409, which leaves a net increase of 10,514 persons. During the last six months of the year the births numbered 11,963, and arrivals by sea 15,354, being a total of 27,317 persons. The deaths for this period were 5,023, and departures by sea 9,684, both together amounting to 14,707 persons. The net increase then will be 12,610 persons.

The number of marriages celebrated during the year was 4,630, being only 25 in excess of the Marriages, number for the previous year. The registered births numbered 23,298, and the deaths 11,193, both showing an increase of 770 and 422 respectively, on the figures for 1875.

Table showing the number of Marriages, Births, and Deaths, registered in the Colony of New South Wales, during each Quarter of the Year 1876.

Quarter ended—	Marriages.		Births,			Deaths.	
What ter ended-	Marriages.	Males.	Females.	Total.	Males.	Females.	Total.
			В				9
Sydney—		440	398	838	405	353	758
31st March	321		401	865	364	335	699
30th June	349	464	460	873	294	271	565
30th September	296	413	1			214	469
31st December	298	377	357	734	255		409
Total	1,264	1,694	1,616	3,310	1,318	1,173	2,491
Suburbs—						1	
31st March	113	343	378	721	245	271	516
	101	389	383	772	265	240	505
30th June			427	832	172	160	332
30th September	117	405			169	163	332
31st December	132	381	352	733	109		332
Total	463	1,518	1,540	3,058	851	834	1,685
Country Districts -							
31st March	832	2,071	2,020	4,091	1,076	662	1,738
30th June	713	2,064	1,984	4,048	1,137	817	1,954
30th September	687	2,299	2,206	4,505	1,113	626	1,739
			2,141	4,286	1,013	573	1,586
31st December	671	2,145	2,141	4,200		3/3	
TOTAL	2,903	8,579	8,351	16,930	4,339	2,678	7,017
New South Wales-	Processing of the Paris of the						
31st March	1,266	2,854	2,796	5,650	1,726	1,286	3,012
30th June	1,163	2,017	2,768	5,685	1,766	1,392	3,158
30th September	1,100	3,117	3,093	6,210	1,579	1,057	2,636
			2,850	5,753	1,437	950	2,387
31st December	1,101	2,903	2,050	3,733		950	-,30/
TOTAL	4,630	11,791	11,507	23,298	6,508	4,685	11,193

MARRIAGES.

From the quarterly table of Marriages we find that of the total number (4,630) celebrated during the year there were 1,264 or 27:30 per cent. in Sydney, 463 or 10 per cent. in the Suburbs, and 2,903 or 62:70 per cent. in the Country Districts. In comparison with the previous year there were more marriages in the City and Suburbs and fewer in the Country Districts.

The ratio of marriages for every 1,000 persons of estimated population in the middle of the year is for the City, 13:55; Suburbs, 6:26; and Country, 6:45. For the whole Colony the ratio per 1,000 is 7:51, the average of the previous nine years being 7:81. Looking at the table which distinguishes the number of marriages celebrated by the various denominations and by the Registrars, we see that the Church of England heads the list with 1,759; Roman Catholic, 921; Presbyterian, 761; Wesleyan, 487; Congregationalist, 232; Primitive Methodist, 101; Registrars, 275; and the balance is made up by other denominations, with double figures only. Those denominations which show an increase on the previous year are as follows:—Church of England, 141, and Congregationalist, 46; Registrars' Offices also show an increase of 25.

The following denominations show a decrease, viz.:—Roman Catholic, 32; Presbyterian, 145; Wesleyan, 15; Primitive Methodist, 8. Of the total number of marriages then the Church of England celebrated 37:97 per cent.; Roman Catholic, 19:89; Presbyterian, 16:44; Wesleyan, 10:52; Congregationalist, 5:01; Primitive Methodist, 2:18; and Registrars, 5:95.

(B.)

Denomination.	1867.	1868.	1869.	1870.	1871.	1872.	1873.	1874.	1875.	1876.
Church of England	832	1,024	1,018	1,159	1,232	1,381	1,547	1,572	1,618	1,759
Roman Catholie	881	912	100	908	885	836	992	837	953	921
Presbyterian	767	698	663	548	648	827	914	886	906	76:
Wesleyan	303	333	313	363	364	397	462	518	502	48
Primitive Methodists	49	51	58	70	60	73	60	60	93	IO
Congregationalist	100	96	106	82	99	148	164	180	186	23:
Baptist	38	21	25	37	30	51	44	52	46	30
Christian Israelite										
Unitarian	2	2	4	3	4	3	3	I	ı	
Hebrew	9	13	13	4	IO	6	10	IO	11	1
Protestant Episcopal						,	2	5	16	I
Free Church of England	186	350	446	451	395					
German Evangelical	3	2	2	7	4	9	7	14	17	I
Christians		I	6	ī	3	3			4	
Free Gospel Church			I							
Free Christian Church									2	
Registrars' Offices	256	233	243	215	219	191	179	208	250	27
TOTAL	3,426	3,736	3,799	3,848	3,953	3,925	4,384	4,343	4,605	4,63

MARRIAGE TABLE of Increase and Decrease.

Denomination.	1875.	1876.	1875. Per cent.	1876. Per cent.	Increase. Per cent.	Decrease. Per cent.
Church of England Roman Catholic Presbyterian Wesleyan Congregational Primitive Methodist Baptist Hebrew Protestant Episcopal German Evangelical Christians, &c. Registrars' Offices	1,618 953 906 502 186 93 46 11 16 17 7 250 4,605	1,759 921 761 487 232 101 39 14 16 19 6 275	35'14 20'69 19'67 10'90 4'04 2'00 1'00 0'23 0'37 0'38 0'15 5'43	37.97 19.89 16.44 10.52 5.01 2.18 0.84 0.30 0.36 0.41 0.13 5.95	2:83 0'97 0'18 0'07 0'03 0'52	0.80 3.53 0.38 0.16 0.01

From the total number (9,260) of persons married during the year 1876, there were 1,516, or 6.65 per cent. who were minors, of whom 132 were males and 1,384 females. Comparing these figures with the returns of the previous year, we find a decrease of 24 persons. Taking the number of minors married in the City, Suburbs, and Country, we find the percentages are 9.77, 13.71, and 19.67, respectively.

TABLE OF Minors.

		No. of Marriages	Min	nors.
		No. of Marriages.	Male.	Female.
Sydney		1,264	22	225
Suburbs Country Districts		463	18 92	109
	TOTAL	4,630	132	1,384

MINORS and Marks.

		Males.	Females.	Total.
Sydney Suburbs Country Districts		2 3 26	13 3 127	15 6
Country Districts	TOTAL	31	143	174

Considering the condition of those who contracted marriage during the year, we find that out of 9,260 persons 4,139 were bachelors, 4,111 spinsters, 489 widowers, and 518 widows. There were also 2 divorced men and 1 divorced woman, who were again married. In the City the bachelors and spinsters were about equal, viz., 1,091 and 1,092 respectively; the numbers in the Suburbs were 409 and 415 respectively, and Country 2,639 and 2,604 respectively. Widowers and widows were nearly equal in the City and Suburbs; in the former 173 and 171, and in the latter 54 and 48 respectively, while the Country shows 262 widowers and 299 widows.

TABLE of Bachelors, Spinsters, &c.

<i>a</i> ,	No. of Marriages.	Bachelors.	Spinsters.	Widowers.	Widows.	Divorced Men.	Divorced Women.
SydneySuburbsCountry Districts	1,264 463 2,903	1,091 409 2,639	1,092 415 2,604	173 54 262	171 48 299	2	I
TOTAL	4,630	4,139	4,111	489	518	2	I

From the total number (9,260) of persons married, we find that 970, viz., 443 males and 527 females, appear as unable to sign their names, so that 10.47 per cent. affixed their marks to their marriage certificates. This shows an improvement of 2 per cent. on the previous year. The percentage of this class to the total number of persons married in the city is 7.63; in the Suburbs 3.56; and in the country 16.71. Out of the 970 persons in the Colony unable to affix their signatures, there were 174 who were minors, namely, 31 males, and 143 females, or 18 per cent. These minors were distributed as follows:—Sydney, 15; in Suburbs, 6; and in the Country Districts, 153.

	No. of	F	ull Signature	es.		Marks.		Marks
	Marriages.	Males.	Females.	Total.	Males.	Females.	Total.	per cent.
Sydney	1,264 463 2,903	1,192 451 2,544	1,143 442 2,518	2,335 893 5,062	7 ² 12 359	121 21 385	193 33 744	7.63 3.26 16.41
	4,630	4,187	4,103	8,290	443	527	970	10.47

(C.)

TABLE showing the number of MARRIAGES registered in the Colony of New South Wales during each quarter of the Year 1867 to 1876.

		Estimated Population at the middle of the Year.	31 March.	30 June.	30 Sept.	31 Dec.	Total.	Ratio per 1,000 living.
Sydney	[1867 1868 1869 1870 1871	68,587 71,310 74,310 77,131 79,540	294 304 332 297 320	277 321 330 313 325	260 281 299 323 316	246 286 314 302 255	1,077 1,192 1,275 1,235 1,216	15.70 16.71 17.15 16.01
syuney	1872 1873 1874 1875 1876	79,745 83,000 86,213 89,796 93,269	305 316 277 324 321	256 346 339 324 349	264 299 306 297 296	230 297 302 287 298	1,055 1,258 1,224 1,232 1,264	13.23 15.15 14.19 13.72 13.55
			3,090	3,180	2,941	2,817	12,028	
	1867 1868 1869	47,429 49,313 51,387	61 68 91	50 72 52	71 61 64	75 52 74	257 253 281	5.42 5.46
Suburbs	1870 1871 1872 1873	53,33 ⁸ 55,004 63,160 65,73 ⁸	72 46 73 90	59 47 71 65	60 60 65 69	52 61 77 92	243 214 286 316	4.55 3.88 4.52 4.80
	1874 1875 1876	68,281 71,119 73,872	80 102 113	81	77 101 117	75 109 132	313 426 463	4·58 5·99 6·26
			796	712	745	799	3,052	
	1867	322,927 335,749	565 582	466 544	55 ² 602	509 563	2,092 2,291	6·47 6·82
Country Districts	1869 1870 1871	349,877 363,162 374,504	574 646 657	546 539 661	572 576 630	551 609 575	2,243 2,370 2,523	6.41 6.52 6.73
	1872 1873 1874	384,777 400,479 415,976	679 729 689	648 710 729	613 685 707	644 686 681	2,584 2,810 2,806	6.44 6.41
	1875	433,264 450,025	786 832	731	740 687	690 671	2,947 2,903	6.80
			6,739	6,287	6,364	6,179	25,569	1
	[1867 1868 1869	43 ⁸ ,943 45 ⁶ ,372 475,574	920 954 997	793 937 928	883 944 935	830 901 939	3,426 3,736 3,799	7.80 8.18 7.98
New South Wales	1870 1871 1872	493,631 509,048 527,682	1,015 1,023 1,057	911 1,033 975	959 1,006 942	963 891 951	3,848 3,953 3,925	7.79 7.76 7.44
	1873 1874 1875 1876	549,217 570,470 594,179 617,166	1,135 1,046 1,212 1,266	1,121 1,149 1,169 1,163	1,053 1,090 1,138 1,100	1,075 1,058 1,086	4,384 4,343 4,605 4,630	7.98 7.61 7.75 7.51
	(20/0	0.77,100	10,625	10,179	10,050	9,795	40,649	/ 51

BIRTHS.

The total number of births registered during the year 1876 was 23,298, consisting of 11,791 males and 11,507 females, the former being 283 in excess of the latter. The percentage of males is 50.61, and of females 49.39. The increase in the number of births on the previous year is as follows:—Males, 411; females, 359; in all, 770. The birth rate of the Colony per 1,000 of estimated population at the middle of the year was 37.75, being a slight decline on the previous year. The average for the previous nine

years was 39.58, so that the birth rate of 1876 was nearly 2 per 1,000 lower than this average. Of the number of births registered in the City the percentage of males was 51.18, and of females 48.82. In the Suburbs female births were in excess of male, the percentage being 50.36 for the former, and 49.64 for the latter. The figures for the Country Districts are for males, 50.67; for females, 49.32. The City birth rate was 35.48, being 1.44 lower than the figures for the previous nine years average. In the Suburbs the average for the previous nine years was 44.52, and the rate of the year 41.39, showing a decline of 3.13. The rate for the Country Districts was 37.62, which is lower by 1.78 than the average of the nine previous years. Comparing the birth rate, per 1,000 of population, of the three divisions with the rate of the previous year, we find the City and Country were nearly stationary, while the Suburbs show an increase of 0.50.

(D.)

TABLE showing the number of BIRTHS registered in the Colony of New South Wales during each Quarter of the Years
1867 to 1876.

		Estimated Population at the middle of the year.	31 March.	30 June.	30 Sept.	31 Dec.	Total.	Ratio per 1,000 living.
Sydney	1867 1868 1869 1870 1871 1872 1873 1874 1875	68,587 71,310 74,310 77,131 79,540 79,745 83,000 86,213 89,796 93,269	591 650 637 685 724 700 760 762 748 838	632 680 762 757 752 729 755 775 858 865	709 678 782 773 802 727 826 824 797 873	676 585 664 694 702 723 724 760 759 734	2,608 2,593 2,845 2,909 2,980 2,879 3,065 3,121 3,162 3,310	38'02 36'36 38'28 37'71 37'48 36'09 36'92 36'20 35'21 35'48
			7,095	7,565	7,791	7,021	29,472	
Suburbs	1867 1868 1869 1870 1871 1872 1873 1874 1875	47,429 49,313 51,387 53,338 55,004 63,160 65,738 68,281 71,119 73,872	534 532 581 596 632 580 696 637 685 721	557 557 679 679 652 612 700 713 740 772	600 631 615 654 666 659 710 729 785 832	598 557 644 617 651 585 657 708 698 733	2,289 2,277 2,519 2,546 2,601 2,436 2,763 2,787 2,908 3,058	48·26 46·17 49·02 47·73 47·28 38·56 42·03 40·80 40·89 41·39
			6,194	6,661	6,881	6,448	26,184	
Country Districts	1867 1868 1869 1870 1871 1872 1873 1874 1875	322,927 335;749 349,877 363,162 374;504 384,777 400,479 415,976 433,264 450,025	3,079 3,364 3,298 3,234 3,521 3,687 3,755 3,824 3,893 4,091	3,079 3,399 3,430 3,493 3,537 3,624 3,509 3,903 4,113 4,048	3,725 3,594 3,679 3,892 3,993 3,941 4,202 4,331 4,282 4,505	3,537 3,258 3,472 3,574 3,601 3,683 4,150 4,212 4,170 4,286	13,420 13,615 13,879 14,193 14,562 14,935 15,616 16,270 16,458 16,930	41.55 40.55 39.66 39.08 38.88 38.81 38.99 39.11 37.98 37.62
			35,746	36,135	40,054	37,943	149,878	
New South Wales	1867 1868 1869 1870 1871 1872 1873 1874 1875	438,943 456,372 475,574 493,631 509,048 527,682 549,217 570,470 594,179 617,166	4,204 4,546 4,516 4,515 4,877 4,967 5,211 5,223 5,326 5,650	4,268 4,636 4,871 4,929 4,941 4,965 4,964 5,391 5,711 5,685	5,034 4,903 5,076 5,319 5,371 5,327 5,738 5,884 5,864 6,210	4,811 4,400 4,780 4,885 4,954 4,991 5,531 5,680 5,627 5,753	18,317 18,485 19,243 19,648 20,143 20,250 21,444 22,178 22,528 23,298	41.73 40.50 40.46 39.80 39.57 38.37 39.04 38.87 37.92 37.75

The number of illegitimate births registered during the year was 950, which gives a percentage of megitimate.

4.07 to total number of births. Taking population into account the rate per 1,000 for the Colony was

1.54, which is somewhat over the rate of the previous year. The percentage to total number of births in

each of the divisions was as follows:—Sydney, 9.24; Suburbs, 2.87; Country, 3.28. A large number of illegitimate births are registered from the Benevolent Asylum; hence the high percentage for the City as compared with the Suburbs and Country Districts:—

(E.)

Table showing the number of Children born out of Wedlock in the Colony of New South Wales, from 1st January, 1867, to 31st December, 1876.

		Sydney.			Suburban.			untry Distri	cts.	Total.			
Year.	Males.	Females.	Total.	Males.	Females.	Total.	Males.	Females.	Total.	Males.	Females.	Total.	
			1.		1								
1867	65	77	142	32	33	65	232	231	463	329	341	670	
1868	67	82	149	30	33 28	58	253	239	492	350	349	699	
1869	106	7.8	184	29	28	57	274	228	502	409	334	743	
1870	90	95	185	52	38	90	250	230	480	392	363	755	
1871	105	IOI	206	30	31	61	259	256	515	394	388	782	
1872	94	. 88	182	34	32	66	287	281	568	415	401	816	
1873		IOI	199	42	44	86	293	311	604	433	456	889	
1874		124	241	40	48	88	305	301	606	462	473	935	
1875		128	260	41	44	85	321	281	602	494	453	947	
1876	170	136	306	43	45	88	309	247	556	522	428	950	
	1,044	1,010	2,054	373	371	744	2,783	2,605	5,388	4,200	3,986	8,186	
								-					

Triplets and Twins. There was one case of triplets recorded during the year, and 195 cases of twins, of which last 9 children were still-born, leaving 381 children, which gives a percentage of 1.63 to total number of births.

(F.)

TABLE showing the number of TRIPLETS and TWINS born in the Colony of New South Wales from 1st January, 1867, to 31st December, 1876.

		Triplets. Twins.								
Year.	No. of Cases.	Males.	Females.	Total.	No. of Cases.	Males.	Females.	Total.		
				-						
867					158	148	166	314		
868	3	7	2	9	162	174	150	324		
869	4	6	6	12	165	161	169	339		
870		I	5	6	188	189	187	370		
871		2	I	3	194	198	190	370		
872	5	7	8	15	179	173	183	35		
373	I		3	3	189	204	173	377		
874	4	9	3	12	177	182	171	353		
875	I		. 3	3	200	180	219	399		
876	I	I	I	2*	195	188	193	381		
	22	33	32	65	1,807	1,797	1,801	3,598		

* One still-bo .

t Two still-born

t One still-born

8 Nine still-born.

DEATHS.

The total number of deaths recorded in the year 1876 was 11,193. In each of the quarters of the year the numbers were as follows:—March, 3,012; June, 3,158; September, 2,636; December, 2,387. Male deaths numbered 6,508, or 58·14 per cent.; and females 4,685, or 41·86 per cent.—the former being 16·82 per cent. higher than the latter. Taking the total number of deaths in the City, we find that the percentage of males is 52·91, and of females 47·09; in the Suburbs, males 50·50, females 49·50; in the Country, males 61·84; females 38·16. The excess of male over female deaths in the City is 5·82 per cent.; in the Suburbs, 1·00 per cent.; and in the Country, 23·68 per cent. The death rate per 1000 of estimated population of the Colony was 18·13, the previous year being 18·12. The average rate for previous nine years was 15·18, so that the rate of the year under report is 2·95 higher than that average. The City death rate for the year was 26·70, which is lower than the figures of the previous year by 3·06, but higher by 3·19 than the previous nine years average. The Suburbs show a rate of 22·81, which is only 0·34 lower than the previous year, and 4·13 in excess of the average for the previous nine years. In the Country districts the rate was 15·59, which is 0·70 over the year preceding, and 2·67 higher than the average of the previous nine years.

(G.)

Table showing the number of Deaths registered in the Colony of New South Wales, during each Quarters of the Years 1867 to 1876.

				/			300 July 1	1 1 1 1 1 1 1
		Estimated population at the middle of each Year.	31 March.	30 June.	30 Sept.	31 Dec.	Total.	Ratio per 1,000, living.
			4.					- N. W.
Sydney	1867 1868 1869 1870 1871 1872 1873 1874 1875	68,587 71,310 74,310 77,131 79,540 79,745 83,000 86,213 89,796 93,269	583 413 371 386 332 477 468 460 900 758	797 410 354 355 362 398 430 401 560 699	330 349 377 337 439 438 408 492 562 565	441 427 409 414 450 497 494 724 651 469	2,151 1,599 1,511 1,492 1,583 1,810 1,800 2,077 2,673 2,491	31°36 22°42 20°33 19°34 19°90 22°69 21°68 24°09 29°76 26°70
			5,148	4,766	4,297	4,976	19,187	
Suburbs	1867 1868 1869 1870 1871 1872 1873 1874 1875	47,429 49,313 51,387 53,338 55,004 63,160 65,738 68,281 71,119 73,872	292 216 240 200 182 244 213 298 533 516	551 247 226 193 169 184 248 291 332 505	243 215 220 187 247 192 249 262 295 332	300 278 245 210 285 295 287 348 487 332	1,386 956 931 790 883 915 997 1,199 1,647 1,685	29'22 19'59 18'11 14'81 16'05 14'48 15'16 17'56 23'15 22'81
			2,934	2,946	2,442	3,067	11,389	
Country Districts	1867 1868 1869 1870 1871 1872 1873 1874 1875	322,927 335,749 349,877 363,162 374,504 384,777 400,479 415,976 433,264 450,025	1,312 1,335 1,216 1,216 988 1,418 1,326 1,350 1,749 1,738	1,226 1,273 1,141 1,122 949 1,134 1,180 1,409 1,879 1,954	1,346 1,027 940 994 969 1,110 1,169 1,309 1,452 1,739	1,210 1,035 952 944 1,035 1,081 1,139 1,308 1,371 1,586	5,094 4,670 4,249 4,276 3,941 4,743 4,814 5,376 6,451 7,017	15'77 13'90 12'14 11'77 10'52 12'32 12'02 12'92 14'89 15'59
			13,648	13,267	12,055	11,661	50,631	
New South Wales	(1867 1868 1869 1870 1871 1872 1873 1874 1875 1876	438,943 456,372 475,574 493,631 509,048 527,682 549,217 570,470 594,179 617,166	2,187 1,964 1,827 1,802 1,502 2,139 2,007 2,108 3,182 3,012	2,574 1,930 1,721 1,670 1,480 1,716 1,858 2,101 2,771 3,158	1,919 1,591 1,537 1,518 1,655 1,740 1,826 2,063 2,309 2,636	1,951 1,740 1,606 1,568 1,770 1,873 1,920 2,380 2,509 2,387	8,631 7,225 6,691 6,558 6,407 7,468 7,611 8,652 10,771 11,193	19.66 15.83 14.06 13.28 12.58 14.15 13.85 15.16 18.12
			21,730	20,979	18,794	19,704	81,207	

INFANTILE MORTALITY.

During the year 1876 there were 4,670 deaths recorded of children under five years of age. These deaths then form 41.72 per cent. of the whole, and are 4.31 lower than the previous year, and 2.31 lower than the average of the previous nine years. The percentage of these deaths to total deaths in each of the divisions is as follows:—Sydney, 41.10; Suburbs, 54.42; Country, 38.89; all of these figures being lower than the previous year by 6.19, 1.01, and 4.31 respectively; and also lower than the nine years average by 5.74, 1.63, and 1.44 respectively. Thus the decrease is much more marked in the City than in either the Suburbs or Country Districts.

The large number of deaths, chiefly of adults, which occur in the Infirmary no doubt causes a lower rate of infantile mortality in the City. Taking each year of the decade it will be seen that infantile mortality has been greater in the suburbs than in the City or Country.

367—B

(H.)
INFANTILE MORTALITY.

	Year. 1867 1868 1869 1870 1871 1872 1873 1874	. , ,	Total	Percentage of total		Sydne	у.		Subur	bs.	Country Districts.			
Val	Year.	Total Deaths of the Year.	eaths of Deaths	Deaths under 5 years to Deaths of the year.	Total Deaths of Year.	Deaths under 5 years.	Percentage of Deaths under 5 years to total Deaths.	Total Deaths of Year.	Deaths under 5 years.	Percentage of Deaths under 5 years to total Deaths.	Total Deaths of Year.	Deaths under 5 years.	Percentage of Deaths under 5 years to total Deaths	
							70.6-	06		66.73	T 004	2.287	46.85	
		8,631	4,574	52.99	2,151	1,262	58.67	1,386	925		5,094	2,387		
	1868	7,225	3,105	42.97	1,599	748	46.77	956	512	53.55	4,670	1,845	39.20	
	1869	6,691	2,973	44.43	1,511	712	47.12	931	553	59.39	4,249	1,708	40.19	
	1870	6,558	2,803	42.74	1,492	673	45.10	790	411	52.02	4,276	1,719	40.50	
	1871	6,407	2,705	42.31	1,583	743	46.93	883	487	55.12	3,941	1,475	37.42	
	1872	7,468	3,203	42.91	1,810	815	45.02	915	503	54.97	4,743	1,885	39.74	
	1873	7,611	2,924	38.42	1,800	719	39 94	997	510	51.12	4,814	1,695	35.62	
	1874	8,652	3,768	43.55	2,077	929	44.72	1,199	672	56.04	5,376	2,167	40.31	
	1875	10,771	4,958	46.03	2,673	1,264	47.29	1,647	913	55.43	6,451	2,781	43'10	
	1876	11,193	4,670	41.72	2,491	1,024	41.10	1,685	917	54.42	7,017	2,729	38.89	
		81,207	35,683		19,187	8,889		11,389	6,403		50,631	20,391		
Mea	an of 10 years	8,120	3,568		1,918	888		1,138	640		5,063	2,039		

Table showing the Births and Deaths of Males and Females registered in each Ward of the City of Sydney, from the 1st January to the 31st December, 1876, distinguishing the Deaths under 5 years of age.

			Di-di-				Des	ths.				
Wards.	Population, Census, 1871.		Births.			Under 5 years.			Above 5 years.			
		Males.	Females.	Total.	Males.	Females.	Total.	Males.	Females.	Total.		
r.—Gipps	8,969	201	194	395	53	38	91	76	49	125	216	
2.—Bourke	5,257	75	52	127	19	13	32	246	209	455	487†	
3.—Brisbane	7,269	136	119	255	48	45	93	66	47	113	206	
4.—Macquarie	7,945	127	170	297	44	43	87	55	33	88	175	
5.—Cook	14,694	431	403	834	147	142	289	95	112	207	496	
6.—Fitzroy	12,000	227	227	454	75	77	152	118	101	219	371	
7.—Phillip	8,154	266	226	492	64	75	139	69	71	140	279	
8.—Denison	10,135	231	225	456	79	62	141	64	56	120	261	
Totals	*74,423	1,694	1,616	3,310	529	495	1,024	789	678	1,467	2,491	

*Exclusive of the islands in Port Jackson.

†Deaths in the Sydney Infirmary 300 (197 males and 103 females).

With regard to the health of Sydney it will be seen that the number of deaths in each of the wards of the City (with the exception of Brisbane Ward) was fewer than in the previous year. Comparing the deaths of children under five years of age to the total deaths in each Ward, we have the following results:—Gipps, 42·13; Bourke (in which the Infirmary is situate), 6·57; Brisbane, 45·15; Macquarie, 49·71; Cook, 58·27; Fitzroy, 40·97; Phillip, 49·82; Denison, 54·02. These percentages are all lower than the figures for the previous year, with the exception of Cook Ward, where they are the same.

Looking now to the deaths under five years of age to the total deaths in each of the Suburban districts we find that Balmain heads the list with 63.05 per cent.; Waterloo, 55.34; Concord, 55.00; Paddington, 54.64; Redfern, 54.16; St. George, 53.60; Newtown, 53.55; Glebe, 53.25; St. Leonards, 50.00; and Randwick, 20.68. These figures show an increase in each of the following districts:—Balmain, Newtown, Paddington, Concord, St. George, and St. Leonards; and a decrease in the Glebe, Redfern, Waterloo, and Randwick.

	Percentage of—													
Ward.	Deaths under 5 years to total Deaths in each Ward.	Deaths under 5 years to Births in each Ward.	Total Deaths to Births in each Ward.	Total Deaths to Population in each Ward. (Census 1861 and 1871.)										
	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876.	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876.	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 18										
1. Gipps	62.80 57.70 53.75 51.53 57.05 48.48 45.39 52.34 56.13 42.13	50.99 25.79 22.16 19.30 22.82 22.47 20.96 29.71 37.37 23.04	81.16 44.64 41.23 37.46 40.00 46.34 46.17 56.76 66.58 54.68	3.04 5.13 5.51 1.81 1.22 1.84 1.81 5.38 5.85 5										
	21.71 17.84 13.62 11.41 13.01 13.53 10.74 9.44 12.50 6.57													
3. Brisbane	69.30 55.77 55.40 50.72 58.82 47.91 39.86 55.73 49.05 45.15	55.12 34.25 26.79 24.47 27.30 23.09 18.91 41.99 38.01 36.47	59.52 61.90 48.36 48.25 46.4: 48.16 47.43 74.02 77.49 80.78	3.50 5.44 5.34 5.18 1.84 1.88 5.03 5.89 5.89 5.										
4. Macquarie	67.24 45.26 55.26 46.96 53.02 51.87 46.71 43.24 50.27 49.71	47.85 25.74 28.00 16.48 21.16 25.27 22.45 22.94 33.21 29.29	71.17 56.88 50.66 35.10 39.78 48.71 44.56 53.05 66.07 58.92	2.24 5.10 1.68 1.49 1.32 1.64 1.45 1.45 5.33 5.										
5. Cook	67.77 58.29 55.15 63.72 57.14 55.62 53.77 61.16 58.27 58.27	43.50 30.52 21.61 22.49 18.93 27.51 23.95 30.62 41.00 34.65	64.16 25.37 39.19 35.30 33.12 40.45 44.24 20.07 70.37 59.47	2.05 5.40 5.34 5.60 1.48 5.18 5.52 5.44 3.65 3.										
6. Fitzroy	61.54 54.62 54:61 49.66 52.57 42.75 39.14 47.53 46.91 40.97	40.13 28.41 25.62 25.21 31.56 27.00 23.11 28.72 39.48 33.48	65.21 52.02 46.02 50.77 60.04 63.12 50.03 60.42 84.16 81.72	3.22 3.02 5.24 5.26 5.30 5.12 5.34 3.53										
7. Phillip	71 20 55 81 54 44 59 37 65 62 57 47 53 84 57 14 61 59 49 82	62.24 20.26 30.43 26.76 29.34 27.47 23.09 27.55 38.28 28.25	87.41 52.44 55.90 45.07 44.56 47.80 42.89 48.22 62.15 56.71	4.34 5.18 3.04 5.40 1.36 5.13 5.04 5.69 3.24 3										
8. Denison	73.00 57.14 60.00 54.60 56.27 64.41 54.18 56.56 64.53 54.02	51.33 27.06 24.05 22.13 22.08 39 89 25.82 26.48 44.31 30.92	70.32 48.93 40.09 40.54 40.75 61.94 47.65 46.81 68.68 57.24	3.56 1.69 5.11 5.05 1.69 5.33 5.00 1.62 5.65 5										

* The Infirmary is situated in Bourke Ward.

(J.)

		Percent	age of—	
District.	Deaths under 5 years to total Deaths in each District.	Deaths under 5 years to Births in each District.	Total Deaths to Births in each District.	Total Deaths to Population in each District. (Census 1861 and 1871.)
	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876.	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876	. 1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876.	1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876.
I. Sydney	58.67 46.77 47.12 45.10 46.93 45.02 39.94 44.72 47.29 41.10	48.39 28.84 25.02 23.13 24.93 28.30 23.49 29.76 39.97 30.93	3 82.48 61.66 53.11 51.28 53.12 62.86 58.72 66.55 84.53 75.25	3.78 2.81 2.65 1.93 2.07 2.37 2.36 2.72 3.50 3.27
2. Balmain	69.23 52.08 62.88 52.94 58.06 61.54 57.62 64.62 61.43 63.05	35.04 20.74 19.74 18.30 17.76 22.00 18.88 25.06 33.24 34.13	3 51.18 39.83 31.39 34.57 30.60 35.73 32.77 38.78 54.12 54.13	3 3.50 5.43 5.46 5.28 1.32 1.25 1.41 5.13 3.05 5.04
3. Glebe	. 64.89 56.52 65.67 60.16 52.87 53.93 49 07 57.44 63.52 53.25	51.96 30.11 27.93 24.28 16 91 14.84 14.35 24.93 33.44 28.51	1 80.07 53.28 42.24 40.86 31.98 33.08 35.29 48.62 52.64 53.00	6.06 3.41 3.91 3.31 1.25 1.22 1.29 1.88 5.49 5.42 5.42
4. Newtown	66.11 21.16 24.02 41.77 23.12 47.36 22.41 27.83 22.90 23.22	39.51 55.33 50.43 15.14 10.10 18.12 10.92 53.13 30.54 52.83	3 59'31 43'65 37'82 29'15 35'95 38'32 38'05 40'00 74'19 51'97	7 2.85 2.03 2.05 1.86 1.45 1.67 1.83 2.09 3.69 2.66
5. Redfern	76.00 28.21 68.16 63.12 66.40 62.21 22.12 22.00 60.30 24.16	45.54 24.49 25.49 19.41 23.38 24.35 22.75 23 72 32.00 27.95	5 59.52 41.86 37.40 30.73 35.51 37.17 41.52 41.63 55.50 51.61	5.60 4.03 3.62 3.11 1.98 1.20 5.03 5.51 5.45
*6. Waterloo	66.94 57.14 67.54 68.04 55.34	27.27 24.43 33.44 36.87 32.30	0 40'74 42'76 49'50 54'19 58'35	5 1.91 2.10 2.39 3.08 3.26
7. Paddington	. 57.34 48.76 53.48 45.37 54.60 52.72 44.56 51.04 54.06 54.64	41.80 21.29 22.44 13.00 20.75 24.54 16.14 26.07 32.33 33.48	8 72.95 43.66 41.95 28.64 38.00 46.61 36.14 51.07 59.81 66.06	5 3.93 2.38 2.53 1.58 1.33 1.44 1.31 1.98 2.27 2.77
* Randwick	18.18 40.00 33.33 32.56 50.68	18.18 13.33 17.64 37.03 13.04	4 33,33 23,00 63,04	1 1 04 1.79 1 68
8. Concord	. 54.54 39.13 52.17 36.73 43.90 44.67 50.00 40.10 44.70 55.00	23.62 15.12 18 90 15.04 15.28 16.58 15.13 13.15 23.89 37.42	2 43.30 38.65 36.22 35.25 28.67 37.20 30.26 32.50 53.46 68.10	2 2 3 1 9 1 9 2 2 0 1 1 6 1 3 1 3 1 3 1 4 2 4 3 1 5
g. St. George	64.96 56.36 50.49 55.40 39.80 53.35 37.09 48.43 39.00 53.60	32.84 23.26 10.42 13.89 12.19 12.28 11.16 10.06 21.10 20.80	50.22 45.28 35.28 55.08 30.65 55.45 30.10 33.19 24.13 20.00	2'43 1'94 1'79 1'31 1'16 0'84 1'16 1 19 2'21 2'36
Io. St. Leonards	68.42 47.91 46.94 34.78 45.45 32.72 48.33 44.94 35.90 50.00	33'33 12'63 14'11 8'51 16'21 9'27 14'35 19'51 13'40 25'20	0 48.72 26.37 30.06 24.46 35.67 28.35 30.00 43.41 37.32 50.41	1 2.50 1.30 1.45 1.33 1.58 1.04 1.16 1.43 1.25 5.38

^{*} Waterloo was not created a separate Registry District till January, 1872; Randwick a Sub-district of Paddington in September, 1872; and Waverley Sub-district of Paddington, 1876.

			Population Census, 1871.	Births.	Deaths under 5.	Total Deaths.
1. Sydney	 		76,159	3,310	1,024	2,491
2. Balmain	 		6,886	375	128	203
3. Glebe	 		5,721	319	90	169
4. Newtown	 		7,922	406	113	211
5. Redfern	 		8,092	465	130	240
6. Waterloo	 		6,306	353	114	206
7. Paddington)		(439	147	269
Randwick	 	}	11,411	31	3	23
*Waverley)		(15	3	6
8. Concord	 		3,520	163	61	111
9. St. George	 		5,340	250	67	125
10. St. Leonards	 • • •		5,126	242	61	122

Looking at the table which treats of the different classes of disease which have proved fatal in the Colony during the year, it will be seen that *Zymotic Diseases* have been the cause of 29:53 per cent. of the total number of deaths. This percentage is nearly the same as in the previous year, but 6:07 over the average of the previous nine years. Under the head of *Constitutional Diseases* the percentage is 11:41, which is a shade under the figures of the previous year, and 0:60 lower than the previous nine years' average. *Local Diseases* give a percentage of 35:21, being slightly lower than previous year, and 2:62 under the figures of the average of nine years preceding. Deaths from *Violence* show a percentage of 7:03, as against 6:74 in the previous year.

(K.)

1867. 1868. 1869. 1870. 1871. 1872. 1873. 1874. Per- Per- Per- Per- Per- Per- Per- Per-	1875.	
	Per-	Per- centage.
CLASS I.—ZYMOTIC DISEASES. Order I.—Miasmatic Diseases—Scarlatina, Measles, Diphtheria, Dysentery, Fevers, &c 2.—Enthetic Diseases—Syphilis, Gonorrhea,	27.13	26.36
&c	0.53	0.44
Delirium Tremens, &c	1.66	2.31 0.45
CLASS II.—CONSTITUTIONAL. 35'07 25'49 22'50 18'95 18'38 21'03 17'85 22'38	29.46	29.23
Order I.—Diathetic Diseases—Gout, Dropsy, Cancer, &c. 2.66 2.—Tubercular Diseases—Scrofula, Phthisis, 2.66	3.54	3.28
Hydrocephalus, &c	8.55	7.83
Class III.—Local. 9'57 11'04 11'15 12'53 13'27 13'61 13'05 12'43	11.46	11.41
Order 1.—Nervous—Apoplexy, Paralysis, Insanity, Convulsions, Cephalitis, Brain Disease,		
&c	12.13	11.22
Heart Disease, &c	4.61	4.79
Asthma, Pleurisy, &c	10.13	10.09
nitis, Hernia, &c	6.64	6.36
6.—Generation—Ovarian Dropsy, Uterus	1.62	1.80
7.—Joints—Arthritis, Ostitis, Periostitis, &c. 0.16 0.15 0.15 0.12 0.14 0.12 0.14 0.11 8.—Integumentary—Phlegmon, Ulcer, Skin	0.18	0.18
Disease, &c. 0'28 0'26 0'33 0'21 0'19 0'23 0'06 0'13	0,10	0.50
CLASS IV.—DEVELOPMENTAL. 31.34 35.32 38.41 38.58 41.09 38.44 41.56 40.09	35.62	35.51
Order 1.—Children — Cyanosis, Spina-Bifida, Teething, &c. 4.63 5.15 5.47 5.46 4.62 5.06 4.32 4.36 2.—Adults—Paramenia, Childbirth, &c. 0.03 0.49 0.69 0.70 0.80 0.49 0.74 1.00 3.—Old People—Old Age, &c. 2.49 5.23 4.95 5.32 5.26 5.54 6.42 5.13 4.—Nutrition—Atrophy, Debility 7.38 6.98 7.13 7.37 6.54 6.25 7.06 6.27	4.68 0.98 4.96 5.57	4.23 1.26 5.45 4.98
14.53 17.85 18.24 18.85 17.22 17.34 18.54 16.76	16.10	15.02
Order 1.—Accident or Negligence—Fractures, Contusions, Burns, Drowning, Suffocation,		
2.—Wounds in Battle—Gunshot Wounds, 7. 7. 900 791 702 004	0.12 0.12	6.34
4.—Suicide—Poison, Drowning, Hanging, &c. 0.46 0.55 0.57 0.53 0.44 0.55 0.46 0.65 5.—Execution—Hanging	0.48	0.42
Unclassified—Violent Deaths	0.07	
7.62 8.58 8.45 10.34 9.48 8.80 8.57 7.80	6.24	7.03
Unspecified	0.29	0.90

The following table shows the percentage which the diseases therein named bear to the total number of causes of death registered in Sydney, Suburbs, and Country Districts respectively:—

	Sydney.	Suburbs.	Country.		Sydney.	Suburbs.	Country,
Scarlatina Croup Typhus and Infantile Fever Erysipelas Dysentery Diarrhœa Delirium Tremens and Intemperance. Dropsy Cancer Tabes Mesenterica Phthisis Cephalitis	0.76	14.66 1.90 3.03 0.47 0.70 6.35 1.42 1.25 3.08 6.05 2.14	9°18 1°38 3°89 1°05 1°78 3°12 0°72 1°39 1°38 0°94 4°39 0°87	Apoplexy Paralysis Convulsions Heart Disease, &c. Bronchitis Pneumonia Enteritis Nephria Premature Birth Teething Old age Atrophy and Debility	1.80 1.48 4.49 4.62 4.33 3.29 1.16 1.76 2.41 1.40 2.77 5.22	1.05 0.83 4.98 3.50 4.80 2.85 1.30 1.13 1.36 2.02 3.02 9.67	1.25 1.83 5.45 4.39 3.87 4.27 2.18 0.18 1.63 2.40 6.98 3.76

In the table which now follows is shown the number of deaths arising from the diseases set forth, distinguishing the number of deaths under five years of age:—

	Syd	ney.	Subt	irbs.	Country	Districts.
Zymotic Diseases.	No. of Deaths.	Deaths under 5 years.	No. of Deaths.	Deaths under 5 years.	No. of Deaths.	Deaths unde 5 years.
Manalan					1	-6
Measles	I				34	26
Scarlatina	206	136	247	154	644	385
Diphtheria	17	13	22	15	165	85
Croup		32	32	27	97	87
Whooping Cough	I	I	I	I	15	15
Typhus and Infantile Fever	77	10	51	10	273	66
Erysipelas	19	2	8	I	74	20
Influenza	14	4	5	3	51	23
Metria	11		9		30	
Dysentery	15	2	12	9	125	66
Diarrhœa	133	93	107	95	219	182
Cholera	18	5	23	17	22	12
Rheumatism	II		8		64	I
Privation	I		2		43	
Delirium Tremens and Intemperance	7				51	
	,					
CONSTITUTIONAL DISEASES.						
ronsv	50	II	25	0	98	8
Propsy	48		25 21	9	97	
	•		2 2		24	2
umour	. 4					66
labes Mesenterica	60	- 59	52	52	66	
Phthisis	202	8	102	I	308	I
Iydrocephalus	19	19	6	6	35	35
Local Diseases.						
Cephalitis	44	36	36	33	6 r	36
poplexy	45	*****	26		88	
Paralysis	37		14		129	3
nsanity	3	I	2		39	
Dpilepsy	15		5		42	I
onvulsions	112	IIO	84	81	383	383
Brain disease, &c.	36	4	25	I	64	4
Ieart disease, &c.		- 4	59	ī	308	7
	108		81		272	152
Bronchitis		49		53		
ongestion of the lungs—Pulmonary	19	9	17	9	55	23
neumonia	82	29	48	24	300	99
ung disease, &c	11		6		40	
sthma	3		I		25	
fastritis	22	20	13	10	47	- 33
nteritis	29	17	23	10	153	72
eritonitis	13		8	2	39	5
Iepatitis	31		13		41	2
iver disease	32	3	22		81	3
ephritis	12	3	7	3	15	6
ephria	44	4	19	5	13	
idney disease	II		5	I	46	4
DEVELOPMENTAL DISEASES.						
remature birth	60	60	23	23	115	115
eething	35	35	34	34	169	169
hildbirth	17		12		107	
	69		51		490	
Old age		128	163	163	264	262
trophy and debility	130	120	103	1 103	204	202

The only table which now remains is the following which shows the mean number of deaths for each month of the decade, as also the mean temperature, rainfall, &c.:—

(L)

	Dowie	J		No. of Deaths.	Mean	Mean Height	Ra	in.	Mean Amount
	Period.			Temperatu		Barometer.	Depth.	Days.	of Ozone.
January February			 	194	71.2	29.927	3.76	14	4·6 4·7
March			 	159	70°7	30.071	6.44	15	4'9
April			 	170	64.8	*099	6.67	14	5.5
May			 	167	58.8	120	7.92	15	5.5
June			 	139	55°1	.10I	5.65	II	5.4
July			 	148	52.6	149	3.97	II	5.6
August			 	151	54.7	.096	1.66	9	5.3
September			 	131	58.5	.063	1.08	II	5.6
October			 	136	64.1	*002	2.67	13	5.5
November			 	161	66.6	29.940	3.76	12	5.4
December			 	200	70.6	.910	2.36	10	4.9
Mean	f 12 m	onths	 	159	63.0	30.038	4.48	12	2,1

I have the honor to be,

Sir,

Your obedient servant,

EDWD. GRANT WARD,

Registrar General.

APPENDIX.

A.

Table showing the number of Marriages, Births, and Deaths registered in the Colony of New South Wales, during the year ended 31st December, 1876.

		Births.			Deaths.	
Marriages.	Males.	Females.	Total.	Males.	Females.	Total.
7				1. 9		
. 1,264	1,694	1,616	3,310	1,318	1,173	2,491
. 463	1,518	1,540	3,058	851	834	1,685
					1	
1,727	3,212	3,156	6,368	2,169	2,007	4,176
. 2,903	8,579	8,351	16,930	4,339	2,678	7,017
		1				
4,630	11,791	11,507	23,298	6,508	4,685	11,193
	1,727	Males. 1,264 1,694 463 1,518 1,727 3,212 2,903 8,579	Marriages. Males. Females. 1,264 1,694 1,616 463 1,518 1,540 1,727 3,212 3,156 2,903 8,579 8,351	Marriages. Males. Females. Total.	Marriages. Males. Females. Total. Males.	Marriages. Males. Females. Total. Males. Females. 1,264 1,694 1,616 3,310 1,318 1,173 463 1,518 1,540 3,058 851 834 1,727 3,212 3,156 6,368 2,169 2,007 2,903 8,579 8,351 16,930 4,339 2,678

B.

Table showing the Marriages, Births, and Deaths registered in each District of the Colony of New South Wales, during the year ended 31st December, 1876.

				3					-	-	
			To	tal.		Birt	ths.				
District.	Popula- tion, Census 1871.	Mar- riages.	Births.	Deaths.	Illegit	Including Illegitimate Births.		Illegitimate Births.		Deaths.	
					М.	F.	M.	F.	М.	F.	
I.—North-eastern.											
1. Richmond River	4,528	27	228	49	123	105	3		29	20	179
2. Tweed River	449	I	19	5	12	7			2	3	14
3. Grafton		(8'I	399	100	187	212	I	- 6	64	45	290
McLean	8,965	1 9	113	29	46	67	2	2	18	II	84
4. Macleay River	5	(33	177	66	80	88	2	3	34	32	III
Nambucca River	{ 4,975	3	21	6	II	IO	I		5	I	15
Bellinger River)	(3	33	5	15	18	I	I	4	I	28
5. Port Macquarie.	2,131	12	81	17	42	39		3	14	3	64
6. Manning River	5,171	28	187	- 44	93	94	3	4	31	13	
7. Port Stephens	2,659	14	137	36	81	56	I		23	13	143
8. Dungog	1,950	15	74	II	38	36	2		8		63
g. Clarence Town	1,000	- J	31	12	13	18	2		5	3	19
io. Paterson	2,561	18	III	22	50	61	2	3	16	6	89
I. Raymond Terrace	3,015	21	96	-33	44	52	3		18	15	
12. Maitland	13,670	104	509	207	265	244	8	7	123	84	63 302
3. Morpeth	3,289	22	IIQ	48	61	- 58		2	26	22	0
4. Patrick's Plains	6,699	36	250	84	128	122	4	2	49		71
5. Muswellbrook	3,616	27	114	37	52	62	I I	I	20	35	
t6. Scone	3,793	16	130	35	62	68	I	2	24	II	77
7. Murrurundi	4,143	41	218	82	115	103	5	2		28	95
/. Dialatana	4,143					103	5		54		136
		509	3,047	937	1,527	1,520	42	38	567	370	2,110
IINORTH MIDLAND.						,					
I. Tenterfield	2,349	23	139	24	72	67	2	3	13	II	III
2. Wellingrove	} 2,648	41	123	37	62	61	2		18	19	88
Vegetable Creek)	7	78	21	36	42	4	2	16	5	57
3. Inverell	2,366	59	201	63	99	102	3	5	33	30	138
4. Armidale	9,760	85	454	141	228	226	6	7	81	60	313
		215	995	286	497	498	17	17	161	125	700

B—continued.

	Popula-		Tot	al.		Births	5.		Dea	ths.	Excess of
District.	tion, Census	Mar- riages.	Births.	Deaths.	Including mate B		Illegiti Birt				Births over Deaths.
	1871.				M.	, F.	М.	F.	М.	· F.	
III.—North-western.	6,866 2,068 3,656 2,838 4,289 2,219 3,167 6,709	$ \begin{array}{c} 72\\ 18\\ 5\\ 21\\ 13\\ 41\\ 24\\ 13\\ 38\\ 24\\ 8 \end{array} $	344 118 53 152 89 260 186 210 188 144 45	103 33 14 57 19 84 47 70 89 51 21	182 58 20 69 45 153 92 116 101 70 26	162 60 33 83 44 107 94 94 87 74	7 2 1 4 4 2 2 3 4 2	1 1 3 4 1 3 2 3 6	67 24 9 41 11 60 29 43 51 25 16	36 9 5 16 8 24 18 27 38 26	241 85 39 95 70 176 139 140 99 93 24
10. Narrabri	1,699	26	95 1,884	40 628	983	901	34	1 26	407	9	1,256
TOTAL		303			903	901					-,-3-
IV.—Metropolitan. 1. City of Sydney 2. Balmain 3. Glebe 4. Newtown 5. Redfern 6. Waterloo 7. Paddington Randwick Waverley 8. Concord 9. St. George 10. St. Leonard's	76,159 6,886 5,721 7,922 8,092 6,306 11,411 3,520 5,340 5,126	1,264 85 29 69 32 78 83 4 7 31 18	3,310 375 319 406 465 353 439 31 15 163 250 242	2,491 203 169 211 240 206 269 23 6 111 125 122	1,694 210 141 215 205 178 219 9 4 89 133 115	1,616 165 178 191 260 175 220 22 11 74 117	170 7 4 7 8 4 3 2 5 3	136 2 8 5 8 8 9 1 2 2	1,318 129 89 108 114 88 130 12 4 53 63	1,173 74 80 103 126 118 139 11 2 58 62 61	819 172 150 195 225 147 170 8 9 52 125
TOTAL		1,727	6,368	4,176	3,212	3,156	213	181	2,169	2,007	2,192
V.—EAST MIDLAND. 1. Newcastle	3,083 5,321 11,551 2,886 3,493 1,870 4,032	(171 30 41 12 3 7 36 10 29 48 17 8 9 11	729 265 288 63 16 84 152 117 162 372 109 88 59	399 124 105 14 4 24 106 44 71 318 77 277 31 65 30	403 132 146 35 11 48 68 55 96 182 52 48 31 55 42	326 133 142 28 5 36 84 62 66 190 57 40 28 54	9 2 4 4 5 7 8 IO 2 2 2 3 1 I	10 3.3 2	224 66 55 8 3 12 60 30 40 219 44 254 16 32 15	175 58 50 6 1 12 46 14 31 99 33 23 15 33 15	330 141 183 49 12 60 46 73 91 54 32 28 44
TOTAL		439	2,690	1,689	1,404	1,286	59	45	1,078	611	1,190
VI.—WEST MIDLAND. 1. Carcoar 2. Cowra 3. Hartley 4. Bathurst 5. Sofala 6. Orange 7. Wellington 8. Tambaroora Hill End 9. Mudgee Gulgong 10. Rylstone	. 1,391 . 5,165 . 16,860 . 2,821 . 7,586 . 3,354 3,625	9 64 20 { 13 { 63 40	322 93 197 784 101 400 167 29 118 399 221	64 169 64 17 71 161 78	51 93 392 58 204 80 16 58 185	154 42 104 392 43 196 87 13 60 214 112 46	18 5 7 5 3 12	1 5 12 3 5 6 16 13 1	92 18 46 422 41 104 39 10 37 94 51 20	59 9 41 195 23 65 25 7 34 67 27 20	238
TOTAL		437	2,924	1,346	1,461	1,463	60	62	774	572	1,57
VII.—SOUTH-EASTERN. 1. Wollongong 2. Kiama 3. Berrima Burrawang 4. Shoalhaven 5. Ulladulla 6. Broulee 7. Eden 8. Bega	5,75° 4,529 4,724 1,652 3,729 1,212	28 { 32 20 11 11	1	68 64 24 44 27 36	92 80 4 97 33 55 27	89 102 98 8 98 37 45	4 I 6	2 3 2 2 I	33 2 23 16 22	29 31 21 11	1 12 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
TOTAL		186	-	-	585	596			196	150	

^{*}The Government Lunatic Asylum and Hospital is situated in this district. † The Hospital for the Insane is situated in this district. † The Government Benevolent Asylum is situated in this district.

B—continued.

			To	tal.		Births					Timenega
District.	Popula- tion, Census 1871.	Mar- riages.	Births.	Deaths.	Inclu Illegitima		Illegiti Birt		Dea	ths.	Excess of Births over Deaths.
					М.	F.	М.	F.	· M.	F.	
					/4						
VIII.—South Midland.											
I. Braidwood	11,422	37	275	90	126	149	4	6	44	46	185
2. Cooma	4,383	45	247	79	127	120	2	2	46	33	168
3. Bombala	2,854	38	163	32	87	76	I	2	21	II	131
4. Goulburn	13,756	103	554	266	282	272	16	5	158	108	288
5. Albury	9,190	86	468	197	222	246	I	4	135	62	271
6. Gundagai	4,699	32	216	68	108	108	2	3	41	27	148
Adelong	4,099	26	IIO	34	47	63	2	I	23	II	76
7. Tumut	3,321	16	IIO	29	50	60	2	2	23	6	81
8. Yass	6,600	54	287	124	141	146	5	I	75	49	163
9. Queanbeyan	4,675	40	196	95	105	91	5	I	57	38	IOI
Io. Burrowa	3,865	23	142	. 58	75	67	3	3	38	20	84
Grenfell	3,003	15	105	45	55	50	2		26	19	60
II. Young	5,924	66	323	116	171	152	IO	4	74	42	207
Murrumburrah	3,924	2	33	5	17	16	I	I	3	2	28
TOTAL		583	3,229	1,238	1,613	1,616	56	35	764	474	1,991
IX.—South-western.		1									
I. Deniliquin	2,714	29	137	67	72	65	2	I	48	19	70
2. Moama	901	3	73	27	42	31	I		19	8	46
3. Moulamein	557	I	9	7	4	5			4	3	2
4. Balranald		8	30	21	13	17			16	5	9
5. Bourke	1	27	84	68	45	39	3	2	53	15	16
6. Hay	3,455	44	171	132	89	82	8	2	100	32	39
7. Wentworth		13	46	35	23	23			29	6	II
8. Menindie		24	71	41	33	38	4	I	25	16	30
9. Wagga Wagga	5,903	82	359	149	186	173	3	5	98	51	210
TOTAL		231	980	547	507	473	21	II	392	155	433
GRAND TOTAL FOR NEW SOUTH	503,891	4,630	23,298	11,193	11,791	11,507	522	428	6,508	4,685	12,105
WALES								1		1	1

The Deaths in the District of Liverpool exceed the Births by 189, which will reduce the excess of Births over Deaths to 11,916.

C.
MARRIAGES—1876.

		Sydne	y.		Subi	ırban	Distri	cts.	Co	untry I	istricts	5.		Tot	al.	
Denomination.	Mar-		Mar-		Marks		Mar-		Marks.		Mar-		Marks			
	riages.	М.	F.	Total.	riages.	М.	F.	Total	riages.	М.	F.	Total.	riages.	М.	F.	Total.
Church of England	479	24	35	59	139	9	8	17	1,141	141	157	298	1,759	174	200	374
Roman Catholic	246	10	25	44	55	2	4	6	620	106	106	212	921	127	135	262
Presbyterian	309	20	45	65	82	I	4	5	370	35	37	72	761	56	86	142
Wesleyan	54		I	I	59		I	I	374	37	31	68	487	37	33	70
Primitive Methodist	18	2		2	II		I	I	72	4	4	8	IOI	6	5	I
Congregationalist	79	5	9	14	88		2	2	65		4	4	232	5	15	20
Baptist	21	I		I	4				14	2	2	4	39	3	2	
Hebrew	13				I								14			
German Evangelical	II		2	2					8	I		I	19	I	2	1
Christians	I				3								4			
Protestant Episcopal									16	3	3	6	16	3	3	
Free Christian Church									2				2			
Registrars' Offices	33	I	4	. 5	21		I	I	221	30	41	71	275	31	46	7
TOTAL	1,264	72	121	193	463	12	21	33	2,903	359	385	744	4,630	443	527	97

D 1.

DEATHS from all causes, of Males at different ages, registered in New South Wales, from 1st January to 31st December, 1876.

CLISS I.—ORDER I. F. SMATHER I. F.	200			1 1		1	1	1		шреі	,	1	1	1	1	1	1	1	1	\		d.	
CLASS I.—ORDIDE I. I. SMRIJPON S. MIRIPON 1	CAUSES OF DEATH.	r year.	years.	years.	years.	years.	years.	15 years.	zo years.	25 years.	30 years.	35 years.	40 years.	45 years.	50 years.	55 years.	60 years.	65 years.	70 years.	75	sars and wards.	not specifie	,
CLASS I.—ORDIDE I. I. SMRIJPON S. MIRIPON 1		ndeı	(1)	to 3	to 4	to 5	to ic	to	to	to	40	9	5 to	40	to	to	5 to	o to	5 to	0 to	5 y up	1ges	[ota]
1. Small-proper 1		Ω	н	0	3			of	H'	500	23	30	33	40	4	ĭ	10	9	9	7	7	7	П
18. Checken-pot	CLASS I.—ORDER I.																						
18. Checken-pot	I. Small-pox																						
3. Scarlaina. 27 78 91 79 56 135 36 9 4 3 3 1 2 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	ia. Chicken-pox							1															16
## Section Sec	3. Scarlatina				79	51			9	4			3										
5. Croup. 6. Whosping-cough. 7. Whosping-cough. 8. Whosping-cou	3a. Diphtheria																						
7. Typhus and Infantile 8	5. Croup	12																					
Service Serv	6. Whooping-cough								1 4										-				
Metric	Fever.)		-/			7	-4				-3		-3							0	0		60
10. Carburche 13.	8. Erysipelas															1							/
12. Dysanthery	10. Carbuncle														I				I				2
13. Dispersion	II. Influenza																		A POST OF THE PARTY OF THE PART				
14. Cholers	13. Diarrhea				6						I	2	Í	2	2			4	9		8		253
16. Remittent Fercer	14. Cholera																					-	
18. Pyremia	16. Remittent Fever							I	I							I							
19. Parotitis	17. Rheumatism																1000000						44
Order 2	19. Parotitis																						I
ORDER 2. 1. Styphilis		282	207	161	124	78	204	67	34	40	25	25	39	24	20	24	16	23	34	29	32	5	1,493
I. Syphilis 7 I	0			-		-	-						-			_	-					10	-
2. Sireture of the Urethra 3. Hydrophobis 4. Glanders 7		7	I									I	I	I		3	I						15
Glanders	2. Stricture of the Urethra														-77	3					- 1	-7.5	
Onder 3	3. Hydrophobia																						
Order 3																			-				
I. Privation		7	1												3								
2. Want of Breast Milk. 51	ORDER 3.										1 7	2	2	2	2	2	1	TO	2	т	2	7	12
4. Alcoholism—a. Del. Tremens	2. Want of Breast Milk																						51
a. Del. Tremens		3	,	I			••••	.,,		ĮI.	I		I	2	I	I	I	I	2				15
54 1 2 7 4 16 15 14 13 14 18 9 5 3 9 184	a. Del. Tremens					•••				I	2		3	3	2	2			2			1000	17
Order 4. I. Thrush	b. Intemperance					···					I	2	9	7	9	7	7	7	3	4	I	2	59
1. Thrush 18 1 <t< th=""><th></th><th>54</th><th></th><th>I</th><th></th><th></th><th></th><th></th><th></th><th>2</th><th>7</th><th>4</th><th>16</th><th>15</th><th>14</th><th>13</th><th>14</th><th>18</th><th>9</th><th>5</th><th>3</th><th>9</th><th>184</th></t<>		54		I						2	7	4	16	15	14	13	14	18	9	5	3	9	184
1. Thrush 18 1 <t< td=""><th>ORDER 4.</th><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>	ORDER 4.																						
Total Class I	I. Thrush														1								- 1
Total Class I				Page 1								110 370											
Total Class I		7.8	Т	-								т		т			т						22
CLASS II.—ORDER I. 1. Gout	TI I OL T		-	-								7.1		-									
1. Gout	Total Class 1	301	209	102	124	78	205	-07	34	42	32	31	50	42	37	43	33	44	40	30	37	14	1,733
1. Gout																		1			1		
1. Gout	CLASS II.—ORDER 1.									4													
2. Dropsy 66 2 2 4 2 8 3 I 2 3 3 3 3 8 9 3 10 8 8 8 6 9 100 3. Cancer													I		2	I	I				I		
3a. Tumour 3b. Polypus 3c. Noma 3c.	2. Dropsy	6	2			2	8	3		2	133 69 020	3	3										
3b. Polypus	3 <i>a</i> . Tumour											100000000000000000000000000000000000000	The state of the s		1				The state of the s				
5. Mortification	3b. Polypus					···			-											1832			
ORDER 2. I. Serofula								1	10000									The state of the s	100000000000000000000000000000000000000		The second		
ORDER 2. 1. Scrofula			-						-		-							20	24		-		228
1. Scrofula 4 1 1	0			- 4	4	3		4	3		4			1/	20	20	25	30	24	15	23		
2. Tabes Mesenterica 45 31 4 3 2 1 2 6 12 35 42 54 46 47 43 21 18 19 4 4 4 4 363 363 Hemoptysis		1			I				T	Т	T	/	T	T		Т	Т				1		12
3a. Hæmoptysis	2. Tabes Mesenterica	45	31	4	3		I				4												86
4. Hydrocephalus													-										
Total Class II 77 40 10 10 7 13 10 16 38 47 61 58 65 70 42 44 49 29 19 27 1 733							1		1														
Total Class II 77 40 10 10 7 13 10 16 38 47 61 58 65 70 42 44 49 29 19 27 1 733		71	38	8	6	1	2	6	12	26	12	55	17	18	41	22	10	10	5	1	1	*	405
	M-4-1 Cl- TT	-	-	-	-	-	-	-	-	-		-	-	-	-	-	-		-	1	1		-
	Total Class II	77	40	IO	10	7	13	IO	16	38	47	61	58	65	70	42	44	49	29	19	27	I	733
]			1)	1			1	1		1					1	1 - 1

D 1—continued.

			,		1	,								,	1	1		1	1	1	1	1
CAUSES OF DEATH,	Under r year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to ro years.	to to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c.	23 232 3	17 43	6 I 24 I	10 9	3 5	4	2 I 	2 2 3 4	2 4 4	1 4 4 2 2 1	3552::5:8	1 6 12 2 5 7	I II II 4 I 6	1 9 10 5 3 9	 17 9 1 2 5	 II II 3 2	1 11 9 1 3 6	1 11 19 2 2 8	15 15 2 7	9 18 4 3 	I I	79 111 125 31 1 38 316 77
	258	60	32,	20	9	10	5	II	10	14	28	33	34	37	34	29	31	43	39	36	5	778
ORDER 2. 1. Pericarditis	 1 4				 I	7	2 1 4	 I IO	I 1 6	1 1 14	 I 22	3 17 20	 2 29 31	3 6 29 38	1 5 20 26	; 3 31 35	1 2 31	1 2 35 38	2 22 24	 1 26	2	34 310 355
ORDER 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a Congestion of the Lungs, pulmonary. 5. Asthma 6. Lung Disease, &c.	85 43 12	1 39 18 4	1 10 13 3 1	I 7 I 4 2 I	3 5 4	1 14 2 8 6	1 5 3 I	1 1 6 1	 14 1	I I I I3 2	 7 5 11 1	 5 2 21 1	 5 4 21 6	8 I 22 5 4	 12 1 16 3 3	 9 3 13 2 7	1 17 3 18 2 7 2	 15 1 10 4	28 I IO 	 13 1 2 2	3 2	10 287 26 271 62 19 38
	143	62	28	16	13	31	10	9	16	21	26	32	39	40	35	34	50	39	43	21	5	713
Order 4. I. Gastritis Enteritis Peritonitis Ascites Ulceration of Intestines Hernia Ileus Intussusception Stricture of Intestines. Fistula Stomach Disease, &c. Pancreas Disease, &c. Hepatitis Jaundice Liver Disease, &c. Spleen Disease, &c.	37 6 1 4 2 1 4	12 11 1 	2	 3 		 4 I I 	 I 	I 6 I I I	5 I I I	1 4 2 1	2 2 1 I	1 5 1 2 6 12	2 II 3 II II II 2 5 II II 2	3 	5 1 2 1 2 6 7	2 1 1 1 6 6 7	I 4 2 2 I I 6 I 8 8	4 2 2 I I 4 2 I I I	4 1 1 3 3	 I I I 4		50 103 23 11 58 12 4 3 3 24 48 9 82
	86	27	'2	4	I	6	5	IO	II	II	13	28	29	24,	26	21	26	30	19	8	I	388
Order 5. 1. Nephritis 2. Ischuria 3. Nephria 4. Diabetes 5. Stone 6. Cystitis 7. Kidney Disease, &c	4	2 I	2 2 I	2 I 	2	4 6 2	 I 	 2 I I	3 	 I 	 3 1 1	2	2 3 I 	3 3 1 2	 2 1 1 4	5 1 5	 2 1 2 2 5	 3 1 1	 I 2 7	I I 2 3 2 6		22 1 44 5 7 11 50
	4	3	5	3	2	12	I	5	5	2	5	4	8	9	8	12	12	15	10	15		140
ORDER 7. 1. Arthritis		 I					I I 2				2		I	2	I	 I I 2		 I I				5 2 7 14
ORDER 8. 1. Phlegmon 2. Ulcer 3. Skin Disease, &c	 1 3		 I			 					 I		 I 	1 2 						 I 		2 6 4
Total Class III	500	153	68	43	26	69	30	46	50	64	98	118	143	153	130	133	153	167	135	108	13	2400

D 1-continued

							D	1-c	ontin	nued												
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	10 to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Agesnotspecified.	Total.
CLASS IV.—ORDER 1. 1 Premature Birth 2 Cyanosis 3 Spina Bifida	96 2 5 13 67	 1 45 46		 	 																	96 2 5 14 112
Order 3.						•••										•••	43	48	83	256		430
Order 4. I Atrophy and Debility Total Class IV	269 452	68	4 4	I		I	2 2										43	48	83	256		299 958
CLASS V.—ORDER 1.— ACCIDENT OR NEGLIGENCE. 1 Fractures & Contusions 2 Wounds { a Gunshot b Cuts 3 Burns and Scalds 3a Sunstroke 4 Poison 4a Bite of Snake or Insect 5 Drowning 6 Suffocation 7 Otherwise	2 1 3 2 9 	1 6 2 II 20	6	3 5 4 	2 4 3 1 1	15 2 3 5 2 1 23 1	14 1 1 12 	18 3 1 2 1 1 11 	19 1 2 3 11 	16 3 3 2 4 8 1	18 1 1 1 14 	25 I I I 2 IO 	31 1 2 8 3 	21 1 2 1 12 1 	21 I I 7 I 	14 1 1 1 4 1 	9 1 1 2 4 1 18	11 1 1 1 1 1 	5 2 7	7 2 9	6 5 	264 12 13 42 19 18 4 160 20 2
ORDER 2.—VIOLENT DEATHS IN BATTLE. I Gunshot Wounds 2 Sword and Bayonet Wounds 3 Otherwise												\										
ORDER 3.—Homicide. Murder and Manslaughter	5		···		···		2 -		2			3				···		•••		<i></i>	2	20
Order 4.—Suicide. 1 Wounds { Gunshot Cut, Stab, &c. 2 Poison 3 Drowning Hanging 5 Otherwise					···			 I 	 I	I	4	 I I I 2 I	 I 	1 3 1 1 2	 3 1 3	 I 	 I 	 I	 I 	I		2 12 7 3 12 2
Order 5.—Execution.				-				I	I		4	2		8	7	2	2	· · ·			I	38
Violent Deaths (not classed)	I	20	25		 I2	6 . 1	 I 32	38	 I	 I	 I 41	 I 52	49	 1	 I		20					11
Sudden Deaths (causes unascertained)				,						7									·			
Causes not specified or			1															1				

D 2.

DEATHS from all causes, of Females, at different ages, registered in New South Wales, from 1st January to 31st
December, 1876.

	1	1	1	1		1	1	1	1	1					1) !	, ,		1	P	
CAUSES OF DEATH.	Under 1 year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to ro years.	to to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS I.—ORDER I.										,											1	
I. Small-pox																						7
1a. Chicken-pox	3	2	4	3	2	 I	 I	 I	 I	 I												19
3. Scarlatina	34	84	75	98	58	140	47	17	II	4	7	I	2			I						579
3a. Diphtheria	3	14	14 I	10	8	32	12	2	3		I	I		I			I					102
5. Croup 6. Whooping-cough	7	16	23	IO	6	8														/		74 9
7. Typhus (and Infantile)						.4.												•••				
Fever)	IO	14	7	6 I	6	31	33	17	23	16	19	13	4	5	3	Ι	I 2	5		2		210 41
9. Metria 10. Carbuncle								4	14	14	9	6	I	I					١		I	50
II. Influenza	7	- 4	 I	Ι			 I				2					 I			3	6		28
12. Dysentery	20	7 40	7	3	ш	3 6			2	I	3	3	3	Ι	3	4	5	3 4		8		47 206
14. Cholera	16	3	*			3	I				2			I	I	3				I		31
15. Ague	2	3				I									I							6
17. Rheumatism			I	I		3	3	2	4	5 5	3	4 I	4	Ι	2	I	I	Ι	3	Ι		39
	241	190	-	138	_		101	44	61	49	51	33	17				11	15	7	20		1,457
ORDER 2.			-33								32		-/									-7137
																						-6
Syphilis Stricture of the Urethra	11	I								I				2		I						16
3. Hydrophobia																						
4. Changers																		••••	`	•••		16
0		I						•••						2				•••	•••	•••		
ORDER 3.																		1				
1. Privation	40											I							2		I	4 41
3. Purpura and Scurvy	I										I											2
4. Alcoholism:— a Del. Tremens												I										ı
b Intemperance										I	4	7	7	I	2	2		2	I		,	27
	41	I								I	5	9	7	I	2	2		2	3		I	75
ORDER 4.																						
I. Thrush	12	I																				13
2. Worms, &c			2																	*/* *		2 8
2a. Hydatid	- ""			7		2		I			2	I		I			I					
, , , , , , , , , , , , , , , , , , , ,	12	I	2			2.		I			2	I					I		•••			23
Total, Class I	305	193	135	138	82	231	101	45	61	51	58	43	24	15	14	14	12	17	10	20	2	1571
CLASS II.—ORDER I.																					/	
I. Gout				2	 I		 I			2		 I	2	I	6		9		2	6		I
2. Dropsy 3. Cancer	4	5				4		5	4		3	IO	13	3 9	9	5	6	9	Í	4		73 63
3a. Tumour	I			I						2	2	I	3	3	3	2		I				19
4. Noma		1																				I
5. Mortification	I														I	I		I	I	I		6
V 31	6	7		3	I	5	I	5	4	4	6	12	18	16	19	12	15	14	4	II		163
ORDER 2.															-							
I. Scrofula	4	I				I			I		I				I							9
2. Tabes Mesenterica 3. Phthisis	57	28	6 I	I		3	4	16	24	39	32	39	30	21	13	18	3	 I	2			92
3a. Hæmoptysis	3					3	4	I	-4	39		39					3					249 I
4. Hydrocephalus 5. Abscess	16	II	2									 I										29 I
J. 21000000	80	-	-	I	-	4	-	17	25	-	-	40	30	21	-	18	-		-			
mul Cl II		40	9	-	т	-	4			39	33	-			14		3	I	2			381
Total, Class II	86	47	9	4	I	9	5	22	29	43	39	52	48	37	33	30	18	15	6	II		544
	-																		The same of the same of	-		-

D 2—continued.

																	-						
	CAUSES OF DEATH.	Under 1 year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
	CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c.	23 196 3	II I 28	6	4	2	7 I 2 3	2 I I 3	2 1 6 3	. I 2 2 2 3	 I 2	1 6 2 4 	2 4 1 1 3 2	3 2 4 4 2 8	 5 4 	 6 6 1 2	3 8 2 2 5	 7 3 1 4	3 9	5 10 I 2	3 4 1 		62 48 55 13 2 24 263 48
		222	41	27	14	8	13	8	12	10	3	17	13	23	9	16	20	15	15	18	10	I	515
	ORDER 2. 1. Pericarditis	I 2		 I			2 4	10	1 6 8	7	1 13	18	I II	20	1 1 14 16	16	 1 16 ———————————————————————————————	16	8	6	4		6 4 172 182
1		3											71. 4			-							
The second second	ORDER 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs pulmonary 5. Asthma	64 34	2 28 1 23 5	1 8 6 1	5 4	2 4 2	2 7 1 7 5	 1 3 	 1 7 1	1 4 10 2	1 3 1 2 1	I I II I	 3 8	 2 2 8 I	 5 1	 4 5 	 4 1 10 	7 4 2	 14 1 2	9 5 	 10 3 1		9 174 16 159 29
	6. Lung Disease, &c	I	/I				I	1			I		4	2		3	2		2	•••	I		19
	ORDER 4.	109	60	16	9	8	23	5	10	17	9	15	16	16	6	14	19	13	20	15	16		416
	1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus	30	5 9 I	4	2	2	9	7 1 	2 I	1 4 9 	7 7 1 1	5 3	7 	 4 I	 4 1 1 	2 I I 2	 2 2 I 	2 4 I	5 I	2 	I 2		32 102 37 5 5 3
And the second s	8. Intussusception 9. Stricture of Intestines. 10. Fistula 11. Stomach Disease, &c. 12. Pancreas Disease, &c. 13. Hepatitis 14. Jaundice 15. Liver Disease, &c.	 I 4	 	 I I			 I I	•••		 I I	3	 I 4 2 7	 I 4	I I 2	 I 3	 2 4 2 6	 2 6 3 5	4 5 1	 7 1 2	3 2 4	I I		3 19 37 17 53
	rő. Spleen Disease, &c	 55	15	7	2	2	1 14	10	3	18	21	24	16	20	16	20	21	25	16	12	6	 I	324
	Order 5. 1. Nephritis 2. Ischuria	•••	I	 3 1 		 I	2 2	2 I	I	2	2 I 2	I I	 6 I	1 4 I	 5 1 		I 2	3	I		 I I		12 1 32 3 1 1
		I	2	6		I	4	3	2	2	5	2	8	6	8	I	3	3	3		2		62
	Order 6. 1. Ovarian Dropsy 2. Uterus Disease, &c									 I	1 4	2 3	I 2	2	3	2 I	 I	ı	 I	2	•••		7 20
										I	5	5	3	2	3	3	I	I	I	2			27
	ORDER 7. 1. Arthritis	 I				•••		 I					 I	 I								••••	2 2 2
		I						2					I	I	I								6
	ORDER 8. 1. Phlegmon 2. Ulcer			I	•••				•••				 I		 I	 I	•••	 I 	 I 	•••	I		3 5 3
	Total, Class III	394		58	25	20	60	38	35	55	57	81	70	88	60	71	81	74	64	53	39	2	1543

D 2—continued.

	1		1	.)		1)	1	1)	1	1)	1)	1				1	d.	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to ro years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS IV.—ORDER 1. 1. Premature Birth 2. Cyanosis	102 5 4 5 68	 58		 I 					6 													102 5 4 7 126
Order 2. 1. Paramenia 2. Childbirth (see Metria)								2 8	23	27	I 27	26	I 22									5 136
Order 3.								10	23	28	28	26	23	3			19	24	40	95	2	180
ORDER 4. 1. Atrophy and Debility Total, Class IV		23	5	 I		I			23	28	28	26	23					24	40	95		258
CLASS V.—ORDER I.— ACCIDENT OR NEGLIGENCE. I. Fractures & Contusions 2. Wounds { a Gunshot}	 I 4	1 8 2 7	1 4 2 9 	3 i 5	 6 1 1 	3 I 8 I 2 I I 6	3 2 3 9	I 2 2 5	4 I 5	2 2 2 2 1 7		2 1 1 1 5	2 4	I I 2 2	I I I 3	4 1 1 	1 4 5	 I 	 I 	2 3 I	 I 	23 2 1 55 14 6 30 10
Order 2.—Violent Deaths IN Battle. 1. Gunshot Wounds 2. Sword and Bayonet Wounds 3. Otherwise																						
ORDER 3.—Homicide. Murder and Manslaughter	 I											2	3									7
Order 4.—Suicide. 1. Wounds { Gunshots Cut, Stab, &c. 2. Poison									 I 	 I 	 I 		3	 I 	 I 		 I 					 2 6 1
ORDER 5.—EXECUTION.				1																		
Violent Deaths (not classed)			17	5	8	16	9	5	6	8	2	9		1 6	4	6	6			6		3
Sudden Deaths (cause unascertained)																				1	·	
Causes not specified or ill-defined	I	458	224	173	2	4 321	155		1 175	1 188	208	201	5	I 122	3	1 132	2			171	19	43
	1-1-20	110	1	1 10	0	1	1	1	1	1	1	1	1	100	1	1	1	1000		1	1	1

D 3.

Deaths from all causes, of Both Sexes, at different ages, registered in New South Wales, from 1st January to 31st December, 1876.

	1	1	1	,			1	1		1	1	- 1	1		1	1	1	1	1	1	ed.	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS I.—ORDER 1. 1. Small-pox	 I 9	4 4 162 31 36 4 31 2 6 13 19 10 4 4 1	7	3	3		1 83 16 2 1 551 1 2 1 1 8 8 1	2 26 5 333 I 4		2 7 2 7 29 2 14 1 3 2 1 1 7 5			2 2 2 13 4 1 1 4 5 2 2 5 5								I I	1 35 1097 204 12 163 17 401 101 50 2 70 152 459 63 8 15 83 16 1
	523	397	294	262	160	433	168	78	101	74	76	72	41	31	36	27	34	49	36	52	6	2950
ORDER 2. 1. Syphilis	18	2			P					I	 	 I	I I 	3	3 3	2 I 	3	3	2 2	2		31 18 49
ORDER 3. 1. Privation	91 4 	 I 	 I						 I I	3 I	2 I	4 1 4 16	3 2 3 14	2 I 2 IO	3 I	4 1 2 9	10 1	2 2 2 5	3	2 I	8	46 92 17 18 86
ORDER 4. 1. Thrush		2	1 2			 I 2				8	9 3	 I	 I									32 3 11
Total Class I	30	2	2	262		3	168	79	103	83	89	99	1 66	52	57	47	56	63	46	57.	16	3304
CLASS II.—ORDER 1. 1. Gout	 10 	7 I	297	6 I 	 3 1 4	 12 1 1 15	 4 1	 6 1 1 	6	 5 3 	6 3 3	1 4 16 2 23	1 10 21 3 35	3 12 23 3 I	1 9 19 6 4	1 14 19 2 1 37	2 17 23 2 I	1 17 16 1 3	8 8 1 2	1 15 14 4 34	 I 	11 173 166 30 3 18
Order 2. 1. Scrofula	8 102 3 38	1 59 3 15	 10 3 4 	7 4 1 1	2 4	1 i i 5 7	10	28 I 	2 59 	81 82	I 86 1 88	85 1	78	64 1 	39 2 34 36	36 	22	5 I 6	6	4		21 178 612 4 60 1
Total Class II	163	87	19	14	8	22	15	38	67	90	100	110	113	107	75	74	67	44	25	38	I	1277
	1	1	1	1	1	1	1	-1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

CAUSES OF DEATH.	I year.	years.	years.	4 years.	years.	years.	to 15 years.	20 years.	25 years.	30 years.	5 years.	40 years.	45 years.	50 years.	55 years.	60 years.	65 years.	70 years.	to 75 years.	years and upwards.	Ages not specified.	
	Under	I to z ;	2 to 3 3	3 to 4.3	4 to 5 3	5 to 10	ro to r	15 to 20	20 to 2	25 to 30	30 to 35	35 to 4	40 to 4	45 to 5	50 to 5	55 to 6	60 to 6	65 to 7	70 to 7	75 ye	Ages n	Total.
CLASS III.—ORDER I. 1. Cephalitis. 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions	46	28 I	I2 I	14	5	II I	4 1 2 1	2 2 3 9	3 2 2 6	I 4 4 2 I 4	4 11 7 2 9	3 10 13 3 8	4 13 15 8 	1 14 14 5 3	23 15 2	 14 19 5 	1 18 12 2 3	I 14 28 2 	 20 25 3 	 12 22 5 3	 I 	141 159 180 44 3 62
8. Brain Disease, &c	428	71	45 I	19		5 4	4	7	7		12	9	14	9	6	7	10		9	4	2	579
ORDER 2.	480	101	59	34	17	23	13	23	20	17.	45	46	57	46	50	49	46	58	57	46	6	1293
1. Pericarditis	1 1 6		 I		 I	2 2 11	2 I I4	1 2 16	I I I3	I 2 27	 I 40	3 28	2 49	4 7 43	5 36	1 4 47	1 2 47	1 2 43	 2 28	 1 30	2	17 38 482
	8		I		I	15	17	19	15	30	4I —	32	51	54	42	52	50	46	30	31	2	537
ORDER 3. I. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs, Pulmonary. 5. Asthma.	149 77	3 67 1 41 9	2 18 19 4	I I2 I 8 2	2 7 7 4	3 21 3 15 11	6 6 I	1 2 1 13 2	I 4 24 3	2 4 2 15 3	I 8 6 22 2	 6 5 29 1	7 6 29 7	 8 1 27 6	 16 1 21 3	 13 4 23 	1 24 3 22 2	 29 2 12 4	37 1 15 	²³ I 5 3	3 2	19 461 42 430 91
6. Lung Disease, &c	2	I	I	I -	I	I	I			4	I	6	3	4	6	9	2	10		2		57
ORDER 4.	252	122	44	25	21	54	15	19	33	30	41	48	55	46	49	53	63	59	58	37	5	1129
1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus	67 6 1 5	17 20 1 1 	6	3	3	1 13 1 	I 8 I I 	I 8 2 I I	1 9 10 I	I II 9 2 I 	4 7 4 I	1 6 7 1 	2 3 7 1 2 2	 8 2 2 1 2	7 1 2 4 1 2	 4 2 I I	3 8 2 2 1 2	4 7 3 1 	 6 I I 2	 I I 3	 I 	82 205 60 16 10 11 20
8. Intussusception 9. Stricture of Intestines 10. Fistula	I		 I	 			 I		I		 I		 3			и и 5	I I 4	 I 3	 I 6	 I 		5 6 3 43
12. Pancreas Disease, &c 13. Hepatitis 14. Jaundice 15. Liver Disease, &c 16. Spleen Disease, &c	.7	I I I	I			I I I	 I		 3 1	3 5	6 2 12	6 1 16	7 I 20	5 4 11	10 2 13 1	12 3 12	11 2 13 1	3 13	5 9	5		85 26 135 5
	141	42	9	6	3	20	15	13	29	32	37	44	49	40	46	42	51	46	31	14	2	712
Order 5. I. Nephritis 2. Ischuria 3. Nephria 4. Diabetes 5. Stone 6. Cystitis 7. Kidney Disease, &c.		3	2 5 1 	2 I 	3	6 8	2 I 	3 I I I	5 	 3 2 	1 3 1 	8 1	3 7 I 	4 8 I I 3	I 2 I I 4	1 6 1 7	 5 1 2 2 5	1 4 1 I I I I I I	 I 2 7	1 3 3 6		34 2 76 8 8 12 62
	5	5	II	3	3	16	4	7	7	7	7	12	14	17	9	15	15	18	10	17		202
Order 6. 1. Ovarian Dropsy 2. Uterus Disease, &c										1 4	2 3	1 2	2	3	2 I	 I	I	 I	2			7.
									1	5	5	3	2	3	3		I	I	2			27
ORDER 7. 1. Arthritis	 I 	 I					2 2				2	2	I 	I 2	I	 I I		I I				7 4 9
	ı	I					4				2	2	2	3	I	2		2				20
Order 8. 1. Phlegmon			I I		I	I					 I	 I	 I 	3	 I		 I 	 I		I I 		5 11 7
1	7		2		I	I					I	I		4	I		I	I		2		23
Total, Class III	894	271	126	68	46	129	68	81	5	121	179	188	231	213	201	214	227	231	188	147	15	3943

D 3—continued.

CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	10 to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Agesnotspecified.	Total.
CLASS IV.—ORDER 1. 1. Premature Birth 2. Cyanosis 3. Spina Bifida 4. Other Malformations 5. Teething	198 7 9 18 135	2		 I															.2.			198 7 9 21 238
	367	105		I			•••	•••		•••				`					•••		•••	473
ORDER 2. 1. Paramenia		•••				•••		2 8	23	1 27	1 27	26	I 22	3	•••							136
		•••						10	23	28	28	26	23	3							***	141
ORDER 3. 1. Old Age														•••			62	72	123	351	2	610
ORDER 4. 1. Atrophy and Debility	498	45	9	I		2 .	2	•••			•••	•••						3 = 4-				557
Total, Class IV	865	150	9	2		2	2	10	23	28	28	26	23	3	•••		62	72	123	351	• 2	1781
CLASS V.—ORDER 1.— ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions 2. Wounds { a Gunshot } b Cuts 3. Burns and Scalds 3a. Sunstroke 4. Poison 4a. Bite of Snake or Insect 5. Drowning 6. Suffocation 7. Otherwise	4 1 4 4 3 17 	2 14 4 18 	7 I 9 2 21 2 42	4 8 I I 4	2 10 4 1 1	18 3 3 13 3 2 1 24 1	15 1 3 2 15 	19 3 3 2 1 1 13 	19 1 2 4 3 12 	18 3 5 2 4 10 2	18 1 1 1 14 	25 1 3 2 3 11 	33 1 2 9 4 50	22 I 3 I I 14 I 	22 1 1 1 8 1 	18 1 1 2 4 1 	10 5 2 4 1	II I I I I I I I	5 I 2 	9 5 1 	6 1 5 13	287 14 14 97 33 24 4 190 30 3
ORDER 2.—VIOLENT DEATHS								7			-33	-13	3	15	-00				¥			
IN BATTLE. 1. Gunshot Wounds 2. Sword and Bayonet Wounds. 3. Otherwise	11.3				.1.																	
ORDER 3.—HOMICIDE. Murder & Manslaughter					I	ı	2	***	2	I	2	5	3		I	I				•••	2	27
Order 4.—Suicide. 1. Wounds { Gunshot Cut, Stab, &c. 2. Poison Drowning Hanging Otherwise								 I 	I I 2	I I I	4 1	 I I 2 I	 3 1 	I 4 I I 2	 4 1 3 	 I I 	 2 I	 I 	I		 I	2 14 13 4 12 2
ORDER 5.—EXECUTION.			***		***		***	I	2	3	5		4	9		-2	3	I	1		I	47
Violent Deaths (not classed)						···		4 0 1		I		2			. 0,4	***			•••	1		3
	I I	•••		•••	•••	1		•••	I	I		3	2	2	I	•••		•••	*		•••	14
Total, Class V Sudden Deaths (Cause unascertained.)	40	38	42		20	70	41	43	46	50	43	61	59	54	45	31	26	19	9	16	16	787
				,				-								-						
Causes not specified or ill-defined.	I				3	6	4	I	2	2	3	5	12	4	6	3	5	3			41	101
GRAND TOTALS	2629	948	493	364	237	665	298	252	346	374	442	489	504	433	384	369	443	432	391	609	91	11193

E 1.

DEATHS from all causes of Males at different ages, registered in Sydney, from 1st January to 31st December, 1876.

CAUSES OF DEATH.	Under 1 year.	I to 2 years.	z to 3 years.	3 to 4 years.	4 to 5 years.	5 to ro years.	to to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Agesnotspecified	Total
Cara I Ones																						
CLASS I—ORDER I. Small-pox																						
a. Chicken-pox																						
. Measles							***		I						1							-
. Scarlatina	3	23	21	19	7	18	7	2	2	1	I	.2										IO
a. Diphtheria		2	I	2 I		2																
. Croup	I	5	5	3	2	I																I
. Whooping-cough											:											
Typhus (and Infantile Fever).		3			2	3	4	I	4	I	3	6	2				I					3
B. Erysipelas	I								ī			I				I			I	I		
. Metria				·																		
. Carbuncle													***									
Influenza	I		I	***																		
Dysentery	I 26	17	4	2		I	/ I		I.		I	I	I	I		2	2	1 2	I 2	3	I	6
. Cholera	4	2			I						I		I				I					I
. Ague			:					I			I											
6. Remittent Fever		I	I				I	I														
7. Rheumatism		 I		***				I	I	I	Ι		Ι	I			I					
Parotitis			I																			
	37	55	34	27	12	25	13	6	10	4	8	II	6	4	I	3	5	3	4	4		27
ORDER 2.	3/			-/		-3								-						1		
r. Syphilis	2										I											
2. Stricture of the Urethra													I	I	2	I	I					1
3. Hydrophobia			,																			
. Glanders				•••		***		***	•••												•••	1
	2										I		I	I	2	I	I					
ORDER 3.							1															
r. Privation			***																			2
3. Pupura and Scurvy	21																					
4. Alcoholism—																						
a. Del. Tremens									I	2			I	I		I						1
b. Intemperance											I	5	3	2	3	I	3					
	21								I	2	I	5	5	3	3	3	3				۵.	4
ORDER 4.																						
I. Thrush						1															***	
2. Worms, &c	***										 I		***									1
Tiyaatia			-			-		-							-		-					9
											I											-
Total Class I	60	55	34	27	12	25	13	6	II	6	II	16	12	. 8	5	7	9	3	4	4	I	33
		-			1					-							1					
0 77 0															1.7.							
CLASS II—ORDER 1.														I	I			. I				1.5
. Gout	2	1	I	2	I	I	3			Ι			4	5		4	I			I		3
G. Cancer											I	3	2	I	6	2	6	2	I	I		2
ga. Tumour										I					I							
b. Polypus						I																
Mortification															I							
	2	I		2	° I	2	3			2	2	4	6	7	9	6	7	. 3	I	2		-
ORDER 2.						-	-					-		,			-					
Scrofula	I							I					I									
. Tabes Mesenterica	12	8	3	I	I	I																2
. Phthisis		I	2	I			2	3	14	13	25	18	17	8	4	4	6		2	2		12
ga. Hæmoptysis											I			I								1
. Hydrocephalus	10											- 2										
	23	9	6	2	I	I	2	4	14	13	26	18	18	9	4	4	6		2	2		16
	23	9				_																

E 1—continued.

							1									1			1		rd	1
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total,
CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c.	8 41 3	5	 7	3	2 I	 I I I	 	I I	 	 2 I 	 I 3 2	2 3 I I	3 2 2	 3 2 1 1 	 3 3	 5 1 1 	 2 3 I	2 I	 4 3 	2 		21 29 16 1 8 63 17
	52	14	7	7	3	4	I	2	I	3	6	7	7	7	6	7	7	3	9		•••	155
Order 2. 1. Pericarditis			•				2 2	3	1 2 3	7	9	3	 8 8	1 2 5	7 8	 8 	2	10	 5 5	1 3 4		4 6 72 82
Order 3. 1. Laryngitis	2 17 10 4	7 I 	1 1 4 	 4 1 	 1 4 3 	 3 1 3	 I 			3	3	 I I 	 2 2 6 2 	 4 1 6 	5 2 I	 5 2 2 	I 2 2 I	2 2 I 2	4 2 I	 2 	 I 	3 63 9 48 14 2 6
	33	8	6	5	8	7				4	7	3	12	11	8	9	6	7	7	2		145
ORDER 4. 1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines 10. Fistula 11. Stomach Disease, &c. 12. Pancreas Disease, &c. 13. Hepatitis 14. Jaundice 15. Liver Disease, &c. 16. Spleen Disease, &c.	10 5	3 1						· · · · · · · · · · · · · · · · · · ·	2	2 I	 1 	I	3	I I I I I	 		 2	 I I 	 I 			13 11 9 2 1 1 2 1 7 17 2 16
ORDER 5.	20	5							3	3	4	8		6		9	3	4	2	2		82
1. Nephritis		 	 I 	 	 I /	1 2		 2 I I	I I	 	 2 1 	 2 2	I 2 I	2 2 2	 I	3 	 I I 	 2 1		 		6 22 I I 2 7 7 39
ORDER 7.											-					-						
1. Arthritis		 I					 			····	I			 I								2 2
ORDER 8.								i X				13										
1. Phlegmon														 I 								 I
Total, Class III	-	29	14	13	12	14	5	II	10	18	30	25	42	36	24	38	20	27	24	10	1	508

E 1—continued.

	1)	1	1	1		1		ontin	1	}		1	1		-+	1	1	1	1	d.	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Agesnot specified.	Total.
G TW 0				1						,												
CLASS IV.—ORDER I. I. Premature Birth	28																					28
2. Cyanosis	2																		,			2 I
4. Other Malformations	3																					3
5. Teething	5	7																				12
	39	7																				46
ORDER 3.																	I	5	6	17		29
ORDER 4. 1. Atrophy and Debility	54	5		I		I																61
Total, Class IV	93	12															I	5	6	17		136
CLASS V.—ORDER I.— ACCIDENT OR NEGLIGENCE.																						
I. Fractures & Contusions	I			2		3	I	2	2	3	3	3	5	2	3				2			32
2. Wounds. $\begin{cases} a \text{ Gunshot} \\ b \text{ Cuts} \end{cases}$						2		 I										 I				4
3. Burns and Scalds		2		2	I	3				I			I									10
3a. Sunstroke		1																				
4a. Bite of Snake or Insect																						
5. Drowning	3					I	I	2	2	2				2		Ι	I					12
7. Otherwise																						
	4	2		4	I	9	2	5	4	6	3	3	6	4	3	I	I	I	2			61
ORDER 2.—VIOLENT DEATHS IN BATTLE. I. Gunshot Wounds 2. Sword and Bayonet					\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\																	•••
Wounds.		1																				
3. Otherwise																•••						•••
			J																			
ORDER 3.—Homicide. Murder & Manslaughter	2								I	I	I						1					6
ORDER 4.—SUICIDE.						-								1								
I. Wounds { Gunshot Cut, Stab, &c.										I	 I				2							4
2. Poison								I						I	I							3
3. Drowning												I			I)
5. Otherwise												I]
								I		I	I	2		3	4							12
Order 5.—Execution.												I]
Violent Deaths (not classed) Found Drowned, &c						I	I		I	I	I	I	2	I	,I							I
Total, Class V	7	2		4	I	10	3	6	6	9	6	7	8	8	8	I	2	I	2			91
	-											١,										
Causes not specified or ill- defined.						I				I	I		2	I	3	I	I				16	2'
	290	108	55	49	27	54	26	27	42	49	76	70	88	69	54	57	46	39	39	35	18	1318
		1											1									

E 2.

Deaths from all causes of Females, at different ages, registered in Sydney, from 1st January to 31st December, 1876.

	of F'E			14											_						1	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	to to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years,	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS I.—ORDER 1. 1. Small-pox														/								
2. Measles 3. Scarlatina	4 2	 16 3	I2 2	18 I	 13	27 2	5	2	····	I	I											100 10 2
4. Quinsy	2 I	3 4	9	I	I	I 4	8	4	9	4	6	 5		 I		····						17 1 47
Fever). 8. Erysipelas 9. Metria 10. Carbuncle	2	I		Ί	•••			I	5	I I	2 2	2	I		I			I 	I	I		13
11. Influenza 12. Dysentery 13. Diarrhœa 14. Cholera	2 2 27	 I I5 2	2			I					3	 I 	3	 	1	2	1 4	I	3	6 1 7 1		6 66 8
15. Ague	3									 I 2		 I	2								 	4 2
16. Lyama.	45	45	26	21	15	35	13	7	16	10	15	10	6	2	3	2	5	3	4	16		299
ORDER 2. 1. Syphilis	5									I				I 		I						8
3. Hydrophobia	,																	•••	•••			8
ORDER 3.	5		***	•••			•••	•••				•••		I .	7			•••		***	***	
1. Privation	17				•••	•••													4		I 	1 17
a Del. Tremensb Intemperance										 I	3	1 4	4	 I	 I	 I		2		•••		17
	17	•••			•••					I	3	5	4	I			•••	2		-,-	1	36
ORDER 4. 1. Thrush			 I			 I						 I										 I 2
Total, Class I	67	4.5	I 27	21		36	13	7		12	18	1 16			4				4	16	 I	346
Total, Class I		45	2/		-3	30	-3			12				4		4	5	5	4			340
CLASS II.—ORDER 1. 1. Gout	 I	2				3		 I	I		2	 I		2	 I		 I	I		 I		20
3. Cancer 3a. Tumour 3b. Polypus 4. Noma		 I										I	7	3	3	3	3					23 2 I
5. Mortification	2	3				3		I	ī		2	4	8	5	4	6	4	2	 I	I		48
Order 2.	ı	I							I		I		***									1
2. Tabes Mesenterica 3. Phthisis 3a. Hæmoptysis 4. Hydrocephalus	3 2		2 I					 5 I	 IO	18	8	13	8	5	3	5			I			34 80 1
5. Abscess	27	5	4					6			9	I 1 14	8	5	3	5						1 128

E 2-sontinued.

CAUSES OF DEATH.	Under I year,	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	10 to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy	6	5	2	3	2	5 I			2		 I 2 	 2 I	 2 I	 4 I 	 1 3 	 I 2 I I	 2 I 	3	 1 5 	 I 		23 16 21 2 1
7. Convulsions	30	9	4	3	2	I	I	I	Ι		I	I	6		I		I		2 8	I		19
Order 2.	37	15	6	6	4	8	I	2	3	2	4	4	9	5	7	-	4	5	0	2		138
Pericarditis Aneurism Heart Disease, &c							 I		2	 6	3	 I	3	4	7	7	5	2	 I	 I		1 43
ORDER 3.					1		I	•••	2	7	3		3	4	7		3				•••	44
Laryngitis Bronchitis Pleurisy Pneumonia Congestion of the Lungs—pulmonary.	10 2	1 6 1 5 2	 I 2	 I	2	I 2 I	I	 I	2 4		1 4	I 6	 2 1 3	2	3		3	6	4	5 2	 I	3 45 5 34 5
5. Asthma										I		I			I			I				5
	12	15	3	I	2	5	I	I	6	1	5	8	7	2	5	•••	3	8	6	7		98
Order 4. 1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines 10. Fistula 11. Stomach Disease, &c 12. Pancreas Disease, &c 13. Hepatitis		3 3 1	2				 I		I	I I	2 I I 	 I	 I	 	 I I 		 1 3	2 	 			9 18 4 2 2 2 8 14
14. Jaundice 15. Liver Disease, &c 16. Spleen Disease, &c	2 I					I					3	I	3	I I	I 2		2			 I		5 16
	13	7	2			I	I	I	I	3	9	3	5	3	7	5	9	3	2	5		80
Order 5. 1. Nephritis		I	2			 I 	 	 I 	2	 I 	I	3	3	1 4		 I 	2	I		 I 		6 22 4
		I	2		•••	2	2	I	2	I		4	4	5		2	2	2		2		33
Order 6. 1. Ovarian Dropsy 2. Uterus Disease, &c			***						 I		2	 I			I	 I						6
Oppen s	1			•••		***	***		I		2		•••	I	I	I	•••					7
ORDER 7. 1. Arthritis 1a. Ostitis, Periostitis, &c. 2. Joint Disease, &c														 								I
Order 8. 1. Phlegmon 2. Ulcer 3. Skin Disease, &c								·				ī			;			 I		I		I 2
Total, Class III	62	38	13	7	6	16	6	5	15	14	24	22	28	21	27	21	23	21	17	18		3 404

E 2—continued.

							-		mun	· · · · · ·												
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS IV.—ORDER 1. 1. Premature Birth 2. Cyanosis 3. Spina Bifida	32 3 2 2 10													1								32 3 2 2 23
ORDER 2. 1. Paramenia	49								4		I 2	3										2 17
ORDER 3.							,		4	6	3	3					2	3	16	19		40
ORDER 4. 1. Atrophy and Debility	58	7	3												· · ·			1.			•••	69
CLASS V.—ORDER I.— ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions	107	20	3					I	4	6	3	3					2	3				3
2. Wounds { a Gunshot b Cuts 3. Burns and Scalds 3a.Sunstroke 4. Poison 4a.Bite of Snake or Insect 5. Drowning 6. Suffocation	 I 			 I 	I	 I			I	I		 I I 				I	I	I		 I 	I	12 1
7. Otherwise	4				I	3			I			2								2	I	20
Gunshot Wounds Sword and Bayonet Wounds. Otherwise																						
Order 3.—Homicide. Murder and Manslaughter.	···					*							2									4
Order 4.—Suicide. 1. Wounds— Gunshot Cut, Stab, &c. 2. Poison			,								•••		 I				 I					 I I
3. Drowning													 									2
ORDER 5.—EXECUTION. 1. Hanging	•••	•••		•••																		
Violent Deaths (not Classed) — Found Drowned, &c. Total Class V			•••	,		3			····	2	I	4	3	1			2			2		29
Causes not specified or ill-defined.	ı		•••	• • • • • • • • • • • • • • • • • • • •	2					I			2		3	I	2				16	28
	271	123	47	29	25	59	19	20	48	53	57	63	60	37	41	38	38	32	39	56	18	1173

E 3.

Deaths from all causes of Both Sexes, at different ages, registered in Sydney, from 1st January to 31st December, 1876.

			,)					1								1	1	1		1	pa	
4	ar.					pi	rs.	rs.	rs.	rs.	rs.	rs.	rs.	rs.	rs.	rs.	rs.	rs.	rs.	q	Ages not specified	1 20
CAUSES OF DEATH.	Under I year.	years.	3 years.	to 4 years.	5 years.	to 10 years.	to to 15 years.	20 years	25 years.	30 years.	to 35 years.	to 40 years	to 45 years.	50 years.	55 years.	60 years.	65 years.	70 years.	70 to 75 years	75 years and upwards.	ot si	
	nder	to 2	to 3	to 4	to 5	to ro	to I	to	to	to	to 3	to 4		to	to	to	to	to	to 7	year	yes n	Total.
	D	н	.01	0	4	w	01	15	50	23	30 1	35	40	45	50	55	9	65	70	75	A	Ĕ
CLASS I—ORDER 1.																				-/		
1. Small-pox								****											/			
2. Measles									I													I
3. Scarlatina	7 2	39	33	37	20	45	12	4	3	2	2	2										206
4. Quinsy			I	I																		3
5. Croup	3	8	14	4	3	2																34
7. Typhus (and Infantile	,	7			3	7	12	5	13	5	9	II	2	Ι	Ι		I					77
Fever). 8. Erysipelas	1	I		I					10	-	2	I	-		- т	I		I	2	2		19
9. Metria	3							I	5	I	2	2	I									II
10. Carbuncle																					,	
11. Influenza	3	2	I									2	 I	 I				I	3	6 I		14
13. Diarrhœa	53	32	6	2		2	I		I	I	3	· I	4	I	I	4	6	3	2	10		133
14. Cholera	7.	4			Ι			 I			2 I		I	Ι			I			I		18
16. Remittent Fever		1	I				I	I														4
17. Rheumatism		 I						I	I	2 2	I	I	3	I	 I		I					5
19. Parotitis			I																			I
	82	100	60	48	27	60	-26	13	26	14	23	21	I2	6	4	5	10	6	8	20	I	572
0					-/			-5			-3											
ORDER 2.	7									I	I			I		· I						II
2. Stricture of the Urethra									.5				I	I	2	I	I					6
3. Hydrophobia																						
4. Gianucis																						
	7							*			I		I	2	2	2	I					17
ORDER 3.																					I	I
I. Privation	38													5								38
3. Purpura and Scurvy													I			I						2
4. Alcholism— a. Del. Tremens									I	2		I	I	I	. 6.	I						7
b. Intemperance										I	4	9	7	3	4	2	3	2				35
	38			.,.					I	3	4	10	9	4	4	4	3	2			I	83
															-		-					
ORDER 4.											·											
1. Thrush			I																			I
2a. Hydatid		····				I					I	I										3
			I			I					I	I										4
m.lol T		1	6-	10	0.5	61	26		07	18	20		22	12	10	II	14	8	8	20	2	676
Total Class I	127	100	61	48	27	01	20	13	27	10	29	32		12								
		-						,														
CLASS II—ORDER 1.													-									
I. Gout														I	I			I		2		3
2. Dropsy	3	3	I	2	2	4	3	I	2	I	3	5	5 9	7 4	9	5 5	9	3	2	2 I		50 48
3. Cancer										I		I			I	I				.,.		4
3b. Polypus		т.				 I																2
4. Noma	I	I													I	I				·		3
4	4	4	I	2	2	5	3	I	2	2	4	8	14	12	13	12	II	5	2	3		IIO
		-	-			-	-			-	-		-	-								
ORDER 2.	2	I						I	I		I		I									7
2. Tabes Mesenterica	33	19	5	Ī	I	Ī											6					60
3. Phthisis	3	I	3	I			2	8	24	31	33	31	25	13	7	9			3	2		3
3a. Hæmoptysis 4. Hydrocephalus	12	5	2																			19
5. Abscess												I										I
	50	26	10	2	I	I	2	10	25	31	35	32	26	14	7	9	6		3	2		292
m + 1 Cl TT		- 20	II	1	3	6	5	II	27	33	39	40	40	26	20	21	17	5	5	5		402
Total Class II	54	30	11	4	3		3		1	33	39	1	1				1					
			1 -		,		-								-							

,										,						,			1	1	1	7 1	
	CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
	CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c.	14 71 4	10 1 18	2 II	6 7	4	6 I I 2 2	I	I I 2	 2 2	 2 I I I	2 2 3 3	4 3 2 2	 3 4 1 8	7 3 i i i	 4 3 2 4	6 3 1 2 1	 4 4 1	 4 4 	 5 8 4	2 I I		44 45 37 3 1 15 112 36
		89	29	13	13	7	12	2	4	4	5	10	II	16	12	13	13	II	8	17	4		293
	Order 2. 1. Pericarditis 2. Aneurism 3. Heart Disease, &c	/a,						2 1 3	 1 3	1 4	 1 13	12	4	11	I 2 9	 1 14 15	 1 15	7	12	6	 4 		4 7 115 126
	Order 3.			•••		•••		3					+			-3		-					
	1. Laryngitis	2 27 12 4	1 13 1 6 2	1 2 6 	4 1 1	3 4 3	5 1 1 4	2	 I	 2 4 	3	3 1 7	 2 7 	4 3 9 2	 4 1 8 	 8 3 I	5 2 2	 4 2 	8 2 I	8 4	7 I	 I 	6 108 14 82 19
	6. Lung Disease, &c	····									2		2	I		I		I	3	1	,		11
		45	23	9	6	10	12	2	I	6	5	12	II,	19	13	13	9	9	15	13	9	I	243
	ORDER 4. 1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines 10. Fistula 11. Stomach Disease, &c. 12. Pancreas Disease, &c. 13. Hepatitis 14. Jaundice 15. Liver Disease, &c.	14 11 1 1 2 	6 4 I	2				 I 	 i i 	 2 I I 	I 2 2 2	2 2 2	I · · · · · · · · · · · · · · · · ·	 4 4 8	I I I I I 2 I I 2	I I 3 I 2		 I I 3	2 I I I I I I	 I I 	I		22 29 13 4 2 3 1 15 31 7 32
	16. Spleen Disease, &c											4	4				3	4		*			
		33	12	2			I	I	2	4	6	13	11	16	9	- 8	14	12	7	4	7		162
	ORDER 5. 1. Nephritis		2	3	 	1	3	I	3 I I	3 	2	I 2 I	5	5 	6		1 4 	3 	3 		 I I		12 44 1 1 3 11
		• • • •	2	3	I	I	5	2	5	5	2	4	8	8	7	I	6	4	5	I	2		72
	ORDER 6. 1. Ovarian Dropsy 2. Uterus Disease, &c									 I		2	I		 I		 I						6
	Order 7. 1. Arthritis		 I					I				I			I I 2	· · · · · · · · · · · · · · · · · · ·							3 2
	ORDER 8. 1. Phlegmon												 I		 I				 I		I		I 3
	Total, Class III	167	67	27	20	18	30	11	16	25	32	54	47	70	57	51	59	43	48	41	28		912

E 3—continued.

	- 1		1	-	1	-		,		- 1	- 1			3	1				1	-		
CAUSES OF DEATH.	Under I year.	I to 2 years.	z to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	rs to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and 'upwards.	Ages not specified.	Total.
CLASS IV.—ORDER 1. 1. Premature Birth	60 5 3 5 15						: 															60 5 3 5 35
	88	20																				108
ORDER 2. 1. Paramenia					2			.f I	4 4	5 6	1 2	3	 I	 I								2 17
ORDER 3.																						
I. Old Age																	3	8	22	36		69
ORDER 4. 1. Atrophy and Debility		12	3	I		2																130
Total Class IV	200	32	3	Ī		2		I	4	6	3	3	I	I			3	8	22	36		326
CLASS V.—ORDER 1.— ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions 2. Wounds { a Gunshot} b Cuts 3. Burns and Scalds 3a.Sunstroke 4. Poison 4a.Bite of Snake or Insect 5. Drowning 6. Suffocation 7. Otherwise	2 I 5	2		3	2	5 2 4 I	I	2 I 2	2 I 2	3 2 2 1	3	3	5 	2 2 	3	 I I	 I I	 I I 	2	 I 	 ,I 	35 4 22 1 12 6 1
	8	2		5	2	12	2	5	5	8	- 3	5	6	4	3	2	2	2	2	2	I	81
Order 2.— Violent Deaths in Battle. 1. Gunshot Wounds 2. Sword and Bayonet Wounds. 3. Otherwise																						·
ORDER 3.—Homicide. Murder and Manslaughter	3								I	I	2		2				I					10
ORDER 4.—SUICIDE. 1. Wounds {Gunshot Cut, Stab,&c. 2. Poison Drowning								 I 		I	1	 I I	 I 	 I I 	 2 I I		I					1 5 4 2 1
								I		I	I	2	I	3	4		I					14
ORDER 5.—EXECUTION. 1. Hanging												I					*					1
Violent Deaths not classed (found drowned, &c.)	I					I	I		I	Ī	I	3	2	2	I	'						14
Total Class V	12	2		5	2	13	3	6	7	II	7	II	II	9	8	2	4	2	2	2	Ī	120
Causes not specified or ill-	I				2	Ī				2	I		4	I	6	2	3				32	55
uenneu.	561	231	102	78	52	113	45	47	90	102	133	133	148	106	95	95	84	71	78	91	36	2491
			6	1	-	1			1	1		1		-	1	1	-	1	1	1	1	1

F 1.

DEATHS from all causes of Males, at different ages, registered in the Suburban Districts, from 1st January to 31st December, 1876.

					1				1		1.	1	1	1	1 -	1	1	1	1		ed.	
*	ear.	10	υż	pi	ţ.i	LT3.	ars.	years.	years.	ars.	years.	ars.	ears.	sars.	ears.	ears.	ears.	ears.	ears.	nd S.	Agesnotspecified	
CAUSES OF DEATH.	r ı year.	years.	3 years.	4 years.	5 years.	10 years.	15 years.	20	25 ye	30 years.	35 ye	40 years.	45 years.	50 years.	55 years.	60 years.	65 years.	70 years.	75 years.	ars a	nots	
	Under	to 3	2 to 3	3 to 4	4 to 5	5 to I	to to	IS to	20 to	25 to	30 to	35 to	to to	t5 to	50 to	55 to	60 to	65 to	70 to	75 years and upwards.	Ages	Total.
		-		(1)	1				1	1.4	1		1	1 4]	1	1					
	,																					
CLASS I.—ORDER I.																						
ı. Small-pox				.A.																		
2. Measles		22		17	13	22		 I		2											 I	119
3a. Diphtheria	7 2	3	19	I	I	32 I	5															10
4. Quinsy	I	5	2	4	2	3									,							17
6. Whooping-cough 7. Typhus (and Infantile					 I		2		2	 I		 I	2									
Fever.)	,	I		I	1	3	2	3	2	1	3	1	2		•••		•				•••	19
8. Erysipelas	I								•••				I			I	I			I		5
10. Carbuncle	*																					
II. Influenza	3		I			 I								•••	 I							3 7
13. Diarrhœa	36	14	I	2	I						т.				I	I		3	I	I		61
15. Ague	4	3					,			I		-::						••••				
16. Remittent Fever							I		 I							 I						
18. Pyæmia												Ι				4			/·			3
	55	49	25	25	18	40	9	4	3	4	3	2	4	I	2	3	I	3	I	3	I	256
		-					,			74	-			-	1							
ORDER 2.																						
2. Stricture of the Urethra	2													 I							1.0	2 2
3. Hydrophobia			,																			
4. Опинсон																						
	2													I				I				4
Order 3.													*									
I. Privation										I										I		2
2. Want of Breast Milk 3. Purpura and Scurvy	12		I							*								I				12
4. Alcholism—																						
b Intemperance												I	I									2
	14		I										I			3		ı		I		20
			-									7										
ORDER 4.							, 1									\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \						
1. Thrush	I	I																				2
2a. Hydatid														/								
	I	I																				2
Total Class 1	72	50	26	25	18	40	9	4	3	5	3	3	5	2	2	3		5		4		282
		7									-											
CLASS II.—ORDER 1.		X																				
2. Dropsy																 I	Î I	I	 I	2		2
3. Cancer												2	I	I	I		I	I		I		8
3b. Polypus																	I				• • • • • • • • • • • • • • • • • • • •	
4. Noma					2					•••												ø ;··
				-		141						•••				•••						I
	1	,	I	I								3	I	I	I	I	4	2	I	4		21
ORDER 2.																						
I. Scrofula												*			I							I
2. Tabes Mesenterica 3. Phthisis	10	9	I			 I								6								20
3a. Hæmoptysis									٠٠٠			5	5		4	I	3	I		I		54
4. Hydrocephalus	2										.,.	•••							h. /.			2
	TO	IO	I			I			6	IO	IO	5	5	6	5	I	3	I		ı		77
	12		3 (1)			Section 1		STATE OF			200		J	THE REAL PROPERTY.		No. of Concession, Name of Street, or other Persons, Name of Street, or ot	0					
Total Class II	13	10	2	I		I			6	10	10	8	6	7	6		7	3		5		98

		1	1				1				1	1	-	1			1	1		1	1 75	1
CAUSES OF DEATH.	Under I year.	I to 2 years.	z to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Agesnotspecified	Total.
CLASS III.—ORDER 1.		6																				22
2. Apoplexy	9		2	5		I				 I	Ι		Ι	2	2		I		2	2		23
3. Paralysis										I		I			I			2	I	3		9 2
4. Insanity														2								- 2
6. Epilepsy								I	2							I						4
7. Convulsions	28	8	4	I	2	2		 I			 I			3				.2				45 12
	37	14	6	6	2			2	2		2	2		7	4	3	I	4	4	5	I	108
Order 2.		-4				3																
I. Pericarditis														I								I
2. Aneurism																	I					26
3. Heart Disease, &c						2			I	I	2	2			4			5	4	4		20
						2			I	I	2	2		I	4		1	5	4	4		27
ORDER 3.																				1		
I. Laryngitis	16	12	 I	2		3				I					 I		3	 I	2	 I	 I	43
3. Pleurisy											2							I				3
4. Pneumonia	5 3	4	2 2	I		2	 I		I		I	Ι	3	2	3			2			 I	25 11
Lungs-pulmonary.							-													1		I
5. Asthma													I	Ι		1						2
	24	16	5	4		5	I		I	I	3	I	5	3	4	I	4	4	2	I	2	87
ORDER 4.													1									
I. Gastritis	I																I 2					2 II
2. Enteritis	2	I	I			I					I				I		I					3
4. Ascites			,				I			4		I	I					I				4
5. Ulceration of Intestines 6. Hernia	Ι																•••					
7. Ileus	2														I							3
8. Intussusception 9. Stricture of Intestines																			•••			
10. Fistula																						2
11. Stomach Disease, &c 12. Pancreas Disease, &c											.,.		Ι									
13. Hepatitis															I	I		I		I	1	4
14. Jaundice									 I			2	2					2	 I			12
16. Spleen Disease, &c																						•••
	. 8	I	I			I	I	I	2		2	3	4	I	4	I	5	4	2	I		42
ORDER 5.																						
I. Nephritis		I	I	I		2								2								7
3. Nephria			I	I	I	4	I		I		I		2.,		I	I						12
4. Diabetes														•••								
6. Cystitis											٠						I					I
7. Kidney Disease, &c	•	I												I				I				4
		2	2	2	I	6	I		I		I			3	2	I	I	1				24
ORDER 6.																						
1. Ovarian Dropsy 2. Uterus Disease, &c								1				F										
		1				1															١	
Onnwa t			-						-	-				-	-	-						
ORDER 7.											٠										·	
Ia. Ostitis, Periostitis, &c														 I						1		
2. Joint Disease, &c							-	-	-	-	-				-					7		
													***				-					
ORDER 8.																	·					
2. Ulcer								١					I									I
3. Skin Disease, &c					•••																	
		_		1									I									2
	I					-	-	3	7	4	10	8	71	16	18	6	12	18	12	II	3	291

F 1--continued.

		1	- 1		1	1	1	1	1		1			-	-	1	1				fled	
CAUSES OF DEATH.	Under I year.	I to 2 years.	z to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS IV.—ORDER 1. í. Premature Birth	12 2 2 6																 					12 2 2 12
5. Icecaning	22	6			****										,							28
				(
ORDER 3.						2000											2	6	6	11		25
ORDER 4. 1. Atrophy and Debility	81	4	2.											•••								87
Total Class IV	103	10	2			.11											2	6	6	II		140
CLASS V.—ORDER 1. ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions 2. Wounds {a Gunshot} b Cuts 3. Burns and Scalds 3a. Sunstroke	24.		i	I		2	I	ī	 I		 I 	3	2	I			I		I	I		15 1 2 1
4. Poison	 I 		 	3	I	5	I	2	 		 I	I		I ,						 		14 1
Order 2.			1	4	2	7	3	3		,	1	4		2					1			
VIOLENT DEATHS IN BATTLE. 1. Gunshot wounds 2. Sword and Bayonet Wounds. 3. Otherwise																						
																,.						inin.
ORDER 3.—HOMICIDE. Murder and Manslaughter	•••															•••	,,,					/* /
ORDER 4.—SUICIDE. 1. Wounds { Gunshot Cut, Stab, &c. } 2. Poison 3. Drowning 4. Hanging															λ. 							
5. Otherwise												1								•••		
										-							I		.,.			I
ORDER 5.—EXECUTION. 1. Hanging										*			-	9				-				•••
Violent Deaths (not classed)	-	./.									1.1					,						
Total Class V	I		I	4	2	7	3	3	I		I	4	2	2			2		I	I		35
Causes not specified or ill-defined.					•••			I			I	2			1				•••		I	5
	259	103	45	42	23	65	15	II	17	19	25	25	24	27	26	II	24	32	21	32	5	851

F 2.

Deaths from all causes of Females, at different ages, registered in the Suburban Districts, from 1st January to 31st December, 1876.

			4	- 1		, ,			11061		7	1	1)	, *	1	-	1	1	1	1 ==	1
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
							1		7								1	1				
CLASS I.—ORDER I. 1. Small-pox 1a. Chicken-pox																						
2. Measles 3. Scarlatina 3a. Diphtheria 4. Quinsy	11	2I 3	16 I	20 I	8 2	34 4	9 I	5	2	 I 	I											128
5. Croup6. Whooping-cough7. Typhus (and Infantile	2 I	3 1	5	2 2	I 3	2 3		6	5		2	 I	 									15 1 32
Fever.) 8. Erysipelas 9. Metria 10. Carbuncle			:43						I	4	4							2				3 9
11. Influenza 12. Dysentery 13. Diarrhœa	1 2 32	2					I		 I	;;;=		 I			 I	 I		 		 I		5 46
14. Cholera 15. Ague 16. Remittent Fever 17. Rheumatism	9															2	f		 I			5
18. Pyæmia							I					2 I		#"								1 2
	58	41	22	25	14	43	15	II	II	8	7	5	I		I	3	I	3	I	I		271
0																· ·	.4					1
ORDER 2. 1. Syphilis 2. Stricture of the Urethra	2								٠													12
3. Hydrophobia																						
	2													;							• • • •	2
ORDER 3.																				/		
2. Want of Breast Milk 3. Purpura and Scurvy 4. Alcoholism— a Del. Tremens	9 1																					9 1
b Intemperance			,																			
	10								· · · ·									\		511		10
ORDER 4. 1. Thrush	I								,													I
2a. Hydatid						I	•••															
Marial Class T	I					I		11		8						3	I	3				285
Total Class I	71	41	22	25	14	44	15	1		_	/	5	-			3	-					
CLASS II.—ORDER 1. 1. Gout	2	3	\	 I				2						Ι	2			 I		 I	l	1 16
3. Cancer 3a. Tumour		I				I						I	2 I	2	I	2	I	I		I		13
3b. Polypus																						
4. Noma 5. Mortification		:::																				
	2	4	/···	I		I		2				I	3	3	3	2	5	2		2		31
Order 2.	I																					1
2. Tabes Mesenterica 3. Phthisis	18	10		I			1	I	2	7	IO	IO	4	5	2	6						32 48
4. Hydrocephalus	3	I		т			т		2	7	10	10	···	5	2	6			•••			85
Total Class II	24	15	3	2		I	I	3	2	7	10	10	7	8	5	8	5	2		2		116
2000																						

F 2—continued.

CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS III—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsey 7. Convulsions 8. Brain Disease, &c.	8 29 I		3 4		2	2 I I	 I I	 I 			2	 I 	 I I 	 	3	 I 3		2 3	 I 	 2 I 		13 13 5 1 1 39 13
ORDER 2. 1. Pericarditis	38		7	· · · · · · · · · · · · · · · · · · ·		1 2	3	2	 I	2	3	2	5		2		5	 I	 I	 I		85 I 33 34
ORDER 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs—pulmonary. 5. Asthma	 10 8 2	1 6 2 1	 4 1	I	 	I 2 I		 I I	2	 I I I	, 	 I 	 I	 I 	 I	2 I	 I 	3	 3 2 	 4 1		2 38 4 23 6
ORDER 4. 1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites	8 4	IO I 2	5	2	2	4	 1	2		3		 2 I			 2 I	3		4	5	5		77 11 12 5 2
5. Ulceration of Intestines 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines 10. Fistula 11. Stomach Disease, &c 12. Pancreas Disease													 I I									I I
13. Hepatitis 14. Jaundice 15. Liver Disease, &c. 16. Spleen Disease, &c. ORDER 5.	 I	3					I		I	 I 5	I I 3	 	1 2 6	3	1 1 1 	2 2 1 8	1 2 4	2				9 4 10
1. Nephritis 2. Ischuria 3. Nephria 4. Diabetes 5. Stone 6. Cystitis 7. Kidney Disease, &c.			 I 		 I 	 I 	· · · · · · · · · · · · · · · · · · ·					3	 I 									 7 ,
Order 6. 1. Ovarian Dropsy 2. Uterus Disease, &c					 					 I	 I	3 						 I	2	*::		2 4
ORDER 7. 1. Arthritis	···						 I						 I 									 I I
ORDER 8. 1. Phlegmon	 I						7														 I	I I 269

F 2-continued.

CAUSES OF DEATH.	Under I year.	r to z years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to '55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS IV.—ORDER 1. 1. Premature Birth 2. Cyañosis 3. Spina Bifida 4. Other Malformations 5. Teething	11 1 2 12																					II I 2 22
	26	10							-11					·						·/		36
ORDER 2. 1. Paramenia								I	3	3	2	 I										1 12
								I	3	3	2	I	3				.2.					13
ORDER 3. 1. Old Age																	5	2	4	15		26
ORDER 4. 1. Atrophy and Debility	66	8	2									***	·					·				76
Total Class IV	92	18	2					I	3	3	2	I	3				5	2	4	15		151
CLASS V.—ORDER I.— ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions 2. Wounds— a Gunshot		1.	1																			Ι
b Cuts 3. Burns and Scalds 3a.Sunstroke 4. Poison		2		 I 			I					f				 						4 1
4a.Bite of Snake or Insect 5. Drowning 6. Suffocation 7. Otherwise	2																	 7				2
	2	2	I	I			2															8
ORDER 2.— VIOLENT DEATHS IN BATTLE. 1. Gunshot Wounds 2. Sword and Bayonet Wounds. 3. Otherwise																						
																		.4			•••	
Order 3.—Homicide. Murder and Manslaughter.									4			·			,			•				
ORDER 4.—SUICIDE. 1. Wounds— Gunshot Cut, Stab, &c. 2. Poison																						 I
3. Drowning																						
Order 5.—Execution.									I						,							
Violent Deaths (not				-,.																		
classed.) Total Class V	2	2	I	I			2		I													9
Sudden Deaths (cause un- ascertained.)			'									\							, <u>,</u>			
Causes not specified or ill- defined.			1					I						I	ā.						I	4
	262	91	42	31	19	57	25	21	23	29	28	28	28	17	17	27	23	20	16	28	2	834

F 3.

Deaths from all Causes of Both Sexes, at different ages, registered in the Suburban Districts, from 1st January to 31st December, 1876.

	V 1:									er, J							- 1					
CAUSES OF DEATH.	Under r year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to IS years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS I.—ORDER 1. 1. Small-pox		43 6 8 I 2 3 23 4		37 2 6 3 2 2	 21 3 3 4 		 14 2 5 	9		 3 4 4 	I 4 I I		3 I				 	2 4	 	I 1 2	 I 	247 22 32 1 51 8 9 5 12 107 23 8 2
	113	90	47	50	32	83	24	15	14	12	10	7	5	I	3	6	2	6	2	4	I	527
ORDER 2. 1. Syphilis 2. Stricture of the Urethra 3. Hydrophobia 4. Glanders	4													I				 I 				4 2
ORDER 3. 1. Privation 2. Want of Breast Milk 3. Purpura and Scurvy 3. Alcoholism— a. Delirium Tremens b. Intemperance	21 3 		 I				'			 		 I 	 I					 I 		 		2 21 5 2
ORDER 4. 1. Thrush	2 2	1 91 ···		50		 I I 84												8				3 4 567
CLASS II—ORDER 1. 1. Gout 2. Dropsy 3. Cancer 3a. Tumour 3b. Polypus 4. Noma 5. Mortification	3	3 1	 	2 2		 I 		2 2			:::	3 4	 3 I 	3 	2 2 4	 1 2 	1 5 2 1 	2 2 	 I 	 3 2 1		3 25 21 2 1
Order 2. 1. Scrofula 2. Tabes Mesenterica 3. Phthisis 3a. Hemoptysis 4. Hydrocephalus	28 5	 19 1 1	4	 I 		 I 	 I 	 I 	 8 	 17 	20	 15 	9	 II 	1 6 7	 7 7	 3 	 I 		 I 		2 52 102 6
Total, Class II	37	25	5	3		2	I	3	8	17	20	19	13	15	II	10	12		77			214

F 3--continued.

CAUSES OF DEATH.	Under 1 year.	I to 2 years.	z to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS III.—ORDER 1. 1. Cephalitis	17 57	6 10	5 8	5 2		3 3	 I 	 I I 	 2 I	 I I 	 3 	 I I 	 2 I 	 3 2 	 5 1 	 2 I 4	 I 	4 5	 3 2 	4 4		36 26 14 2 1 5 84 25
	75	16	13	7	4	7	2	3	3	2	4	4	3	8	7	7	2	9	6	9	2	*193
ORDER 2. 1. Pericarditis	 I					1 4	3	2	2	3	 5 5	4	 5 5	I I	6	 I	6	6	 5 5	5 5		2 59 61
Order 3. I. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs—pulmonary. 5. Asthma 6. Lung Disease, &c	 26 13 5	1 18 6 1	 5 3 2	2 I	I	 4 2 3	 I	 I I 	 2 I 	I I I 	 2 I 	I I I	 3 2	 3 1	 1 4 	2 I	1 4 1 	4 I 2 I	5 2	5 I	 I I	4 81 7 48 17
	44_	26	10	6	2	9	-I	2	3	4	3	3	7	5	6	4	6	8	7	6	2	164
Order 4. 1. Gastritis	9 6 2 I 2 	3 3	 I 			 I 	I	 I 	2 I I	2 I I	2	I I I I I I I I I I I I I I I I I I	 I I I I 4	I	I I I	2 I	2 2 1 	I	I			13 23 8 6 2 4 1 2 13 4 22
	21	4	I			1	2	I	4	5	5	4	10	4	8	9	9	6	3	I		98
ORDER 5. 1. Nephritis		I I	I 2 3	I I 2	2	2 5 7	 I 		 		 I 	3	I	2 I	 I 2	 I 	 I 	 				7 19 1 5
ORDER 6. 1. Ovarian Dropsy 2. Uterus Disease, &c										 I	I	I	77.					 I	2			2 4
2. Cterus Disease, ac										I	I) I						I	2			6
ORDER 7. 1. Arthritis							 I						 I	 I								 I 2
ORDER 8. 1. Phlegmon 2. Ulcer 3. Skin Disease, &c	2												I									 I 2
Total, Class III	7.12	48	27	15	8	29	10	8	13	15	19	19	28	23	30	22	24	31	23	21	4	560

F 3-continued.

		-					1							-			1	1		1	G.	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	to to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS IV.—ORDER 1. 1. Premature Birth	23 1 2 4 18																			/ 		23 1 2 4 34
	48	16	/																		•••	64
ORDER 2. 1. Paramenia								I	3	3	2	 I	3									1 12
ORDER 3.													'			,	7	8	10	26		51
ORDER 4. 1. Atrophy and Debility	147	12	4											1	V							163
Total, Class IV	195	28	4					I	3	3	2	1	3				7	8	10	26		291
CLASS V.—ORDER 1.— ACCIDENT OR NEGLIGENCE. 1. Fractures and Contusions 2. Wounds { a Gunshot}		2	2	I I 3	 I 	2	I I I	I	 I 		 I 	3	2	I			I		I	/ / 		16 1 2 5 1
	3	2	2	5	2	7	5	3	1		I	4	2	2			I		I			42
ORDER 2.— VIOLENT DEATHS IN BATTLE. 1. Gunshot Wounds 2. Sword and Bayonet Wounds	•••																					
ORDER 3.—HOMICIDE. Murder and Manslaughter						,																
ORDER 4.—SUICIDE. 1. Wounds { Gunshot { Cut, Stab, &c} } 2. Poison					7				I								 I 			···		I I 2
Order 5.—Execution. 1. Hanging																	.,.	.,.				
Violent Deaths (not classed)														<i>,</i>								
Total, Class V	3	2	2	5	2	7	5	3	2		1	4	2	2			2		I	I		44
Causes not specified or ill-defined.			I					2			I	2		I							2	9
	521	194	87	73	42	122	40	32	40	48	53	53	52	43	44	38	47	52	37	60	7	1685

G 1.

Deaths from all causes, of Males at different ages, registered in the Country Districts, from 1st January to 31st December, 1876.

						01	ב טמב	Jece	mbe.	r, 18	570.											
CAUSES OF DEATH.	Under 1 year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years,	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
																			1			
CLASS I.—ORDER I. I. Small-pox Ia. Chicken-pox 2. Measles 3. Scarlatina 3a. Diphtheria	 i 6 i7 io	 2 33 12	3 51 15		 I 3I	 85 21		 I 6	 I 2	 I				 								 15 293 85
4. Quinsy	10 6 8	 10 2 13	14	9 2 8 5	5 8 	2 4 18	3	3	2			 	2								I	7 55 8
Fever). 8. Erysipelas 9. Metria 10. Carbunele	8	I			I				2		6	8	5	6 I 	2	3	4	4	7	6		49
11. Influenza'	35 68 4	2 4 22 2	4 7	2 2	 I	1 2 3	 I		I I	I 2	3 2	5 	I	1 2 3 1	4 4 1	 I I	1 4 2	1 5 4	4 7 4	6 7 4	 I	37 89 125 11
15. Ague	3			I			4	2	2	I	3	2	2	 I	 I 4 I	2	3	5		 I	 I	4 5 34 2
	190	103	102	72	48	139	45	24	27	17	14	26	14	15	2 I	IO	17	28	24	25	3	964
ORDER 2. 1. Syphilis	3	1										I	I	 I	3	I	2	2	2	2		. 10
3. Hydrophobia												·										
	3	I			1					-/-		I	I		4	I	2	2	2	2		20
ORDER 3. 1. Privation	 18 1									2 I	2	3	3	2 I	3	4	10	2 I	I	 	7	40 18 9
a Del. Tremens b Intemperance							>					4	1 4	7	4	6	4	3	4	 I	2	9 41
	19								I	4	3	10	9	11	10	11	15	8	5	2	9	117
ORDER 4. 1. Thrush 2. Worms, &c. 2a. Hydatid	17					 I				ii.			 I			 I						17 1 2
	17					I							I			I						20
Total Class I	229	104	102	72	48	140	45	24	28	21	17	37	25	27	35	23	34	38	31	29	12	1121
CLASS II.—ORDER 1.					150								I	I		I	I			I		5
2. Dropsy	3	I		I		7 I		I I I		2	2 I I	2 I I	4 5 	4 12	3 3 2	5 12 	6 10 1	7 10	5 6 1	6 8	I	61 70 8
4. Noma 5. Mortification							 I							I -0	2		 I	2	I	2	16	10
ORDER 2.	3	I		I	2	8	I	3	I	2	4	4	10	18	10	18	19	19	13	17	I	155
Scrofula Tabes Mesenterica Phthisis Hæmoptysis Hydrocephalus	3 23 10	14 1 		1 2 I	I 2	 I	4	9	15	19	19	23 	25	29	13	13	10 	3 1	2	ī		8 40 187 1
Transcription in in it.	36	19	I	4	3	I	4	9	16	20	19	24	25	29	13	14	10	4	2	I		254
Total Class II	39	20	İ	5	5	9	5	12	17	22	23	28	35	47	23	32	29	23	15	18	I	409
			1									J)				1		1			

G 1—continued.

								co		wew.							1		1	1	1	1-
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	10 to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Agesnotspecified	Total.
Class III.—Order 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c.	6 163	6 26 32	4 I I3 I	2 4 I	I I 2 4	2 I 	1 1 1	1	2 2 3	I I 2 2 2 I	3 3 5 2 2 5	1 4 8 2 4 5 24	1 7 9 4 1 4	1 4 8 2 2 6	 12 8 1 2 1	 5 10 3 1	1 8 6 I 2 5 23	1 9 16 2 2 6 36	 9 11 2 4	5 15 4 3 2	I I I I I I I I I I I I I I I I I I I	35 69 100 28 1 26 208 48
Order 2. 1. Pericarditis	 1 4 5					2 5 7	 4 5	7	 1 3 4	i 6 8	 I II	3 12	2 21	1 4 24 29	1 4 9	1 2 23 26	1 2 28 31	1 2 20 23	 2 13	19	2	6 28 212 246
Order 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs, pulmonary. 5. Asthma 6. Lung Disease, &c.	 52 28 5	1 20 13 4	 8 7 1	1 1 3 1	2 I I I	1 8 1 8 1	1 4 3 	1 1 6 1	 13 1	I I I O 2	 4 2 7 1	 5 1 19 1	 3 2 12 3	 4 14 5 3	6 I II 2 3	 4 1 11 2 6	 13 3 16 2	 12 6 3 1 6	 22 1 8 	 IO I 2 2	 I	5 181 14 198 37 16 30
ORDER 4. 1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines. 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines 10. Fistula 11. Stomach Diseases, &c 12. Pancreas Disease, &c 13. Hepatitis 14. Jaundice	30 4 I I I 2 	38 9 1 	 1 	7 3	5 I	3 i	8 I I I	9	15 2 1 1 1 	16 4 	16 2 I I I	1 4	22 I I I 2 I	26 3 1 2 4	23 4 1 2 1 1 1 4	24 2 I I I	40 2 1 2 1 1 5 1	28 4 2 1	34 4 	 1 1 1 		35 81 11 5 4 7 8 2 2 2 3 15
15. Liver Disease, &c 16. Spleen Disease, &c	58	21		4		5	3	8	7	3 8	7	7	5 14	17	6 I 21	5	5 18	22	15	5	 I	54 3 264
ORDER 5. 1. Nephritis		-S	I			I 2			 I 	 I 	 I		I I I	I I I I	 I I I 2	 I I 5	 I I 2 5	 I I I 8	 I 2 6	1 1 2 3 2 6		9 1 10 4 6 8 39
ORDER 7. 1. Arthritis					···	3					I	 I	I	4	I	7 I I	9	 I I	9			3 2 4
ORDER 8. 1. Phlegmon	 I 2		 I 			I					 I			I I						 I 		2 4 3 9
Total Class III	325	91	40	18	HI	38	21	32	34	42	58	85	90	101	88	89	121	122	99	87	9	1601

G 1—continued.

		1	1				-				1	1 5	1	T	ī	1	1	1	1	T	j.	1
CAUSES OF DEATH.	Under z year.	I to 2 years.	z to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 79 years.	70 to 75 years.	75 years and upwards.	Agesnotspecified	Total.
Court IV Cour																1						
CLASS IV—ORDER 1. 1. Premature Birth	56														***							56
2. Cyanosis																						2
4. Other Malformations	8	1	,											\							N	9 88
5. Teething	56	32	•••				•••	•••									•••					
	122	33									٠						-,,-				•••	155
0																						
ORDER 3.								•••					`				40	37	71	228		376
		,									4				*							
ORDER 4. 1. Atrophy and Debility	134	13	2				2															151
Total Class IV		46	2			1	2	. 0									40	37	71	228		682
Criss V. Orang							. 1								7							
CLASS V—ORDER 1— ACCIDENT OR NEGLIGENCE.			1																			
1. Fractures & Contusions 2. Wounds \{ a Gunshot \} b Cnts	Ι	I	5		2	10	I2 I	15	17	13	15	19	24 I	18	18	14	8	11	2	6	6	217
(I					I		2	I				1			,	I					7
3. Burns and Scalds	3	4 2	5	3	2 2	3		1 2	3	2		I	I	2 I		 I	I	I		2		31
4. Poison	2			I				I		4	I	2		I	I	I	2	I			I	18
4a. Bite of Snake or Insect 5. Drowning		11	12			17	I 10	7		6	14	9	8	9	7	3	3		2		5	134
6. Suffocation	5		2		I		,						3	1	I	I	I	I				16
7. Otherwise	12	18	24	5	7	37	24	29	31	31	31	33	38	33	29	21	16	16	4	8	12	459*
					-																-	
ORDER 2.—VIOLENT DEATHS						3																
IN BATTLE.																						
2. Sword and Bayonet																						
Wounds, 3. Otherwise				·																		
			-	-		-																
																-						
		,																				
ORDER 3—HOMICIDE. Murder and Manslaughter	2				I	I	2		I			3			I						2	14
Muluel and Manslaughter	3				-							3						- 1	-	-		
ORDER 4.—SUICIDE.																						
I. Wounds { Gunshot Cut, Stab,&c.														I								1 8
2. Poison											3	I		2	I	 I	 I			I		4
3. Drowning													I									I
4. Hanging									I	I		2		2	2	1		I				IO
g. Ceder race									I	I	3	4	I	5	3	2	I	I	I	I	I	25
						-			-													
Order 5.—Execution.										I		I										2
					-							1										
Violent Deaths (not classed)							.,.					,										
		18	24	5	8	38	26	29	33	33	34	41	39	38	33	23	17	17.	5	9	15	500
Total Class V		-	-	-	1																	
Total Class V	1																	1	1			The Party of the P
Total Class V Causes not specified or ill-defined.					I	I	2		I	۵۰	_ I	2	5	2		I	2	3			5	26

G 2.

Deaths from all causes, of Females at different ages, registered in the Country Districts, from 1st January to 31st December, 1876.

A CONTRACTOR OF THE PROPERTY O										r, 18			())	1	1)	1) 751	1
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
						1	6															
CLASS I.—ORDER 1. 1. Small-pox	 3 19 1 7 6	2 47 8 10 1	 4 47 11 9	 3 60 8 7 4	 2 37 6 4 	I 79 26 5 24	 1 33 11 2 	I 10 2 7	 1 8 3 	 I 2 	 5 1 	 I I 	 2 	 I 4		 I 	 I 					 19 351 80 2 42 7
Fever). 8. Erysipelas 9. Metria 10. Carbuncle 11. Influenza 12. Dysentery 13. Diarrhœa 14. Cholera 15. Ague 16. Remittent Fever 17. Rheumatism 18. Pyæmia	9 4 16 57 4 2	 4 4 16 3	 1 5 	 I 3 3 I I I I	I	3 5 3 1 	 I 2	3 	1 8 1 3	1 9 1 1 4 3	2 3 2 I 1	3 4 I I	2 2	I I	 I I I I 	I I I I I I I I I I I I I I I I I I I	2 I I	2 3 2 I		I	I	25 30 14 36 94 11 2 6 30 7
9	138	104	85	92	53	151	73	26	34	31	20	18	10	9	8	6	5	9	2	3	I	887
ORDER 2. 1. Syphilis 2. Stricture of the Urethra 3. Hydrophobia 4. Glanders	4	I			· · · · · · · · · · · · · · · · · · ·									I						,		6
Y	4	-												I								
Order 3. 1. Privation	 14 	 I 		•••	····						 I	 3	3						2 I			3 15 1
	14	I		. 44	١						2	4	3		I	I			3			29
ORDER 4. 1. Thrush 2. Worms, &c. 2a. Hydatid		I	 I					 I			2			 I		···· ···· ···	 I					12 1 5
Total Class I			86	92	-				24	21							I 6					
CLASS II.—ORDER 1.	10/	107		92	53	151	73	27	34	31	33	22	13	11	9	7	6	9	5	3	I	940
1. Gout. 2. Dropsy. 3. Cancer. 3a. Tumour. 3b. Polypus. 4. Noma. 5. Mortification.	I			I		I	 	2	3	2 2	I I 2	7	I 4 2	1 4 3 	3 5 3 	3 I	4 2	7 1 1 	2 I	 4 3 		 37 27 16 4
	2			2		I	I	2	3	4	4	7	7	8	12	4	6	10	3	8		84
Order 2. 1. Scrofula 2. Tabes Mesenterica 3. Phthisis 3a. Hæmoptysis 4. Hydrocephalus	2 18 11	7 5	I			3 	3	 10	 12 	14	14	16	18		8	7	3	 I	 I			4 26 121 17
Total Class II	31	12	2	• • • •		4	3	10	12	14	14	16	18	11)	9	7	3	I	. I.		•••	168
Total Class II	33	12	2	2		5	4	12	15	18	18	23	25	19	21	II	9	II	4	8		252

G 2-continued.

CAUSES OF DEATH.	Under r year.	r to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to ro years.	ro to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total,
CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c.	9 137 1	6 17 1	1 13 	I 6			2 I I I	 I 5 2	I 2 2 2 I	 I	3 4 3	2 I I I	3 1 4 1 2	 3 	 2 3 1 	 6 1 1	5 2 I 2	I 4	 3 4 1 	 I 2 I 		26 19 29 11 16 175 16
ORDER 2. 1. Pericarditis 2. Aneurism. 3. Heart Disease, &c	147 I I		14 1	7		I 2	5 6	9 1 4 6	6 4	 5	 12	7 1 8	 12 12	3 1 9 11	6 7 7	10 8 	6	5 5 - 5	8 4 -4	4 2		5 3 96
ORDER 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs, Pulmonary. 5. Asthma	24 8	 16 16 2	3 3 I	4 2	I I 2	1 4 1 4 3	3	 1 6 	I 6 2	I 2 I I	 I 7 I	I I 2	 I 5 	2	I 3 2	2 I 9 2	3 3	5 1 2	2 I	 I 		4 91 7 102 18
ORDER 4. I. Gastritis 2. Enteritis 3. Peritonitis	77 6 20	35	8 2	6	4 2	1 14 1 9	4 6	7	9 1 3 8	5	10	6	7	 3 1	7 	16	8	8 3	4	4		10 241 12 72 28
4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines							 I		 I			5 2	i	I	I		 I 		 I	 	 I 	1 2 1 7 3
11. Stomach Disease, &c 12. Pancreas Disease, &c 13. Hepatitis 14. Jaundice 15. Liver Disease, &c 16. Spleen Disease, &c	 I		I			I I I			I I I	2 I	1 2 1 3 	 I 3 	3	 2 2 	I I 3	I I 3 	I I I I I I I I I I I I I I I I I I	 4 1 2 	2 1 4 			11 14 8 27 2
ORDER 5.	29_	5	5			13	-	- 12	15	13	12	12	9	10	9	7-	12		- 9		1.	100
Nephritis Ischuria Nephria Diabetes Stone Cystitis. Kidney Disease, &c.			 I 2			 	 			 I I 	 	 I	 	 I I 	 	 I	 I 	 				6 1 3 3 1 7
	I	I	3			I	I	I		4	I	I	I	2	I	I	I	1				21
ORDER 6. 1. Ovarian Dropsy 2. Uterus Disease, &c										3 4	I I 2	 I	2	2	I I 2		I					4 10
ORDER 7. I. Arthritis Ia. Ostitis, Periostitis, &c Joint Disease, &c	I																					14 I I I
ORDER 8.	·I				 I		I					I										3
Phlegmon Ulcer Skin Disease, &c	2		I		 									I	I		I					3 2 7
Total, Class III		65	32	15	9	32	25	25	34	32	48	37	43	31	33	44	39	30	25	II	I	870

G 2—continued.

		,	1					,										,		,	1 -1	1
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	Io to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified.	Total.
CLASS IV—ORDER 1. 1. Premature Birth	59 1 2 1 46.	 1 35		I																		59 1 2 3 81 146
ORDER 2. 1. Paramenia								7 8	16	19	23	22	18	2								109
ORDER 3.	1	\										•••			1		12	19	20	61	2	114
ORDER 4. 1. Atrophy and Debility Total, Class IV		8 44						8	16		23	22		2			12		20	61		113
CLASS V.—ORDER 1. ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions 2. Wounds { a Gunshot} b Cuts 3. Burns and Scalds 3a. Sunstroke 4. Poison 4a. Bite of Snake or Insect 5. Drowning 6. Suffocation 7. Otherwise	4 1	1 6 2 7 7 16	 1 4 2 9 	I I I	 5 1 1 	1 1 7 1 2 1 I 3	1 2 1 3 7	1 2 2 5	3 1 	2 I 2 5		 I I 	2 I I 4	I I I 2 4	ı	4 	3 		 	2 2 2 		19 2 1 39 12 6 30 5
Order 2.—Violent Deaths IN Battle. I. Gunshot Wounds 2. Sword and Bayonet Wounds. 3. Otherwise	7						:::															
Murder & Manslaughter ORDER 4.—SUICIDE. 1. Wounds {Gunshot Cut, Stab,&c. 2. Poison									 I 	 I 	 I 		 		 I 							3 4 1
Order 5.—Execution.				···					I	I		6	2		I							6
Violent Deaths (not classed) Total Class V	10	16	16	3	7	13	7	5	5	6	 I	5	7	4	4	5	4			4		123
Causes not specified or ill-defined.						4	2		I			I	I								2	11
, ,	683	244	136	113	69	205	III	77	105	106	123	110	108	67	67	67	70	69	55	87	6	2678

G 3.

Deaths from all causes of Both Sexes, at different ages, registered in the Country Districts, from 1st January to 31st December, 1876.

								Dec		,												
CAUSES OF DEATH.	year.	years.	ars.	urs.	ars.	sars.	rears.	rears.	25 years.	years.	rears.	vears.	vears.	50 years.	55 years.	60 years.	years.	70 years.	years.	and ds.	Ages not specified.	
CAUSES OF DEATH.	Under 1	to 2 yes	2 to 3 years.	to 4 years.	to 5 years.	to ro years.	to to 15 years.	5 to 20 years.	20 to 25 y	25 to 30 y	30 to 35 years	35 to 40 years.	40 to 45 years.	15 to 50 3	50 to 55 3	55 to 60 y	60 to 65 years.	65 to 70 3	70 to 75 years.	75 years and upwards.	Agesnot	Total.
		н	- A	6	4	w	Ĥ	н _	ñ	o,	ñ	0	4	4	אט	rv	9	9 1	1	1	7	
CLASS I.—ORDER 1. 1. Small-pox																						
Ia. Chicken-pox	Ι.																					I
2. Measles	9	4	7	3	3	I	I	2	2	2												34
3a. Diphtheria	36	80	98	103	68	164	57	16	5	2	5	2 2	2	 I		I						165
4. Quinsy	2			2		2	2														I	9
5. Croup 6. Whooping-cough	17	3	23	15	12	9		 2.														97
7. Typhus (and Infantile Fever.)	17	22	15	9	3	42	34	19	24	20	17.	13	8	10	5	4	2	7	I	I		273
8. Erysipelas	17	I			2	I		3	3 8	I 9	3	11 4	2 I	2 I	2	3	6	6	7	7	 I	74
10. Carbuncle														I				I				2
11. Influenza	15 51	8	1 4	1 2		5			I	1 2	4	6	1 2	2 4	5	I	1 4	8	7	8	 I	51 125
13. Diarrhœa	125	38	12	3 5	2	8	 I		2	3	2		3	4 I	4 2	2	3	6	4	4		219
14. Cholera	8	2		2		3	I			 I	I	I	2		I	I	I	I	 I			22
16. Remittent Fever	5	3		2											I							II
17. Rheumatism				I		6	6	4	5	5	6	3	2	2	6	3	4	6	2	2	I	64
	2		I	I			I			3				A :								9
Oppur	328	207	187	164	101	290	118	50	61	48	43	44	24	24	29	16	22	37	26	28	4	1851
ORDER 2.	7	2										I	I	I	3	- I						16
2. Stricture of the Urethra.					1									I	I		2	2	2	2		10
3. Hydrophobia																						
	7	2				0								2	4	I	2	2	2	2		26
ORDER 3.	/	-																				
I. Privation										2	2	4	3	2	3	4	10	2	3	I	7	43
2. Want of Breast Milk	32	I																				33
3. Purpura and Scurvy 4. Alcoholism—	I								I	I	I	I	I	I	I		I	I		.,.		10
a Del. Tremens						2						2	I	I	2	I		2				9
b Intemperance										I	2	7	7	7	5	7	4	3	5	I	2	51
	33	I							I	4	5	14	12	II	II	12	15	8	8	2	9	146
ORDER 4.									1						1							
I. Thrush	28	I																				29
2. Worms, &c			I			I					2			I			 I			•••		2
zur Hjuntiu	28	I	I			I	-	I	-	300	2	-	I	I	-	I	I	-	-			38
Total, Class I,		-	-	-64		-	8	51	62	52	50	59	38	38	44	30	40	47	36	32		2,061
Total, Class I	390	-		104	-	-9-	-	32		32	30	39	30	30	77	3-	40	47	30	32	-3	
	-							1														
CLASS II.—ORDER I.													I	I		I	I			I		5
2. Dropsy	4	1	1	2	I	8	I	3	4	4	3	2	5	5	6	8	IO	14	7	IO		98
3. Cancer	 I			,		I		I		2	3	8	9 2	16	8 5	12 1	12 I	II	6	11	I	97 24
3b. Polypus											3			3								-4
4. Noma					I		т							 I			 I		2			I
5. Mortification							I							1	3		-	3	2	3		14
	5	I		3	2	9	2	5	4	6	8	II	17	26	22	22	25	29	16	25	I	239
ORDER 2.																				1		
I. Scrofula	5			I		I			I	I		I			I	I		*				12
2. Tabes Mesenterica 3. Phthisis	41	2 I	I	2	I	4	7	10	27	33	33	39	43	40	21	20	13	4	3			308
3a. Hæmoptysis								19				39	43				-3	I				300
4. Hydrocephalus	21	9	2	I	2																	35
	67	31	3	4	3	5	7	19	28	34	33	40	43	40	22	21	13	5	3	I		422
Total, Class II	72	32	3	7	5	14	9	24	32	40	41	51	60	66	44	43	38	34	19	26	I	661
Total, Class 11	72	32	3	1	3	14	9	1 -4	32	1	41] 31			44	43	30	34	19	20		[00]

G 3—continued.

							4	0 0	contin	incu.											(77	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	10 to 15 years.	15 to 20 years.	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS III.—ORDER 1. 1. Cephalitis 2. Apoplexy 3. Paralysis 4. Insanity 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease,, &c	15 300 I	12 43	5 I 26 I	3 10	i	2 I I	3 I 2 I 2	3 7 4	3 2 4 4	1 1 2 2 3 	4 6 5 2 6 8	3 5 9 3 5 6	4 8 10 8 2 6	1 4 11 2 2 6	 14 11 2 2	 6 16 4 1	I I3 8 2 2 7	I IO 20 2 2 6	 12 15 3 	 6 17 5 3 	I I I	61 88 129 39 1 42 383 64
	316	56	33	14	6	4	9	16	13	10	31	31	38	26	30	29	33	41	34	33	4	807
ORDER 2. 1. Pericarditis	1 5 7		 I		 I	1 2 7	10 I	I II I3	 7 8	1 11 13	 I 23	1 3 20	33	2 5 33 40	1 4 16	3 31 35	1 2 34 37	1 2 25 28	 2 17	21	2	31 308 350
ORDER 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia 4a. Congestion of the Lungs, Pulmonary. 5. Asthma 6. Lung Disease, &c.	 96 52 13	1 36 29 6 1	I II IO 2 I	1 5 5 1	3 3 I	2 12 2 12 4 1	1 4 6 1	I I I I I I I I I I I I I I I I I I I	 1 19 3 1	1 3 1 11 3 2	5 3 14 2 I I	6 2 2 I I I 3	3 3 17 3 2 1	4 16 5 3	7 I 14 2 2 4	 6 2 20 4 8	 16 3 19 2 7 1	17 1 8 3 1 6 36	 24 1 9 4 	 II I 3 2	I I 2	9 272 21 300 55 25 40
	163	73	25	13	9	33	12	10	24	21	20	34	29	20	30	40	40	30	30			
Order 4. 1. Gastritis 2. Enteritis 3. Peritonitis 4. Ascites 5. Ulceration of Intestines 6. Hernia 7. Ileus 8. Intussusception 9. Stricture of Intestines 10. Fistula 11. Stomach Disease, &c 12. Pancreas Disease, &c 13. Hepatitis 14. Jaundice 15. Liver Disease, &c 16. Spleen Disease, &c	21 50 4 2 1 1 3 3 2	10 13 1 	 3 	2 3 	3	I I2 I 2 I I	7 I I I I	I 6 I	1 5 9 1 I 1 1 2 I	1 8 6 2 4	2 3 2 I 3 I 6	3	2 3 2 1 2 1 2 2 2 	6 1 2 1 2 4 3 6 1	6 1 1 3 1 2 5 9 1	4 1 1 1 1 1 1 1 1 1 1 1 8	1 6 1 2 2 1 1 1 6 2 6 1	4 5 2 I 2 7 2 10	5 1 2 4 4 8	 I I	 I	47 153 39 6 6 8 15 2 5 3 26 41 15 81
	87	26	6	6	3	18	II	10	22	21	19	29	23	27	30	19	30	33	24	6	2	452
ORDER 5. 1. Nephritis 2. Isehuria 3. Nephria 4. Diabetes 5. Stone 6. Cystitis 7. Kidney Disease, &c	•••	 I	I I 3			2 2	i	 	 I 	 I 2 2	 I 	, i	2 I I I	I 2 I I I	I I I 2	 I 6	 2 1 2 5	 1 1 9	 I 2 6	1 1 2 3 2 6		15 2 13 7 7 7 8 46
	5	I	5	•••		.4	I	2	I	5	2	I	5	6	6	8	10	12	9	15		98
ORDER 6. 1. Ovarian Dropsy 2. Uterus Disease, &c	•••									1 3 4	I I 2	 I	2	2	I I 	•••	 I					4 10 14
ORDER 7. 1. Arthritis 1a. Ostitis, Periostitis, &c. 2. Joint Disease, &c	 I						I	1			1 	2	I		I	 I		 I				4 3 5
ORDER 8. 1. Phlegmon 2. Ulcer 3. Skin Disease, &c	1		I I		ı	I					 			 I 2	 I		 I					4 7 5
Total, Class III	5 584	156	72	33	I 20	70	46	57	68	74	106	122	133	3 132	1 121	133	160	152	124	98	10	16 2471

			,		1					iuea.											· -:	
CAUSES OF DEATH.	Under I year.	I to 2 years.	2 to 3 years.	3 to 4 years.	4 to 5 years.	5 to 10 years.	10 to 15 years.	15 to 20 years	20 to 25 years.	25 to 30 years.	30 to 35 years.	35 to 40 years.	40 to 45 years.	45 to 50 years.	50 to 55 years.	55 to 60 years.	60 to 65 years.	65 to 70 years.	70 to 75 years.	75 years and upwards.	Ages not specified	Total.
CLASS IV—ORDER I. 1. Premature Birth	115 1 4 9	2	 	 I																		115 1 4 12 169
	231	69		I							***											301
ORDER 2. 1. Paramenia				h				1 7			23	22	1 18	2								2 107
								8	16	19	23	22	19	2								109
ORDER 3.	١														A		52	56	91	289	2	490
ORDER 4. 1. Atrophy and Debility		21	2				2								•••	<i></i>			•••		•••	264
Total Class IV	470	90	2	I			2	8	16	19	23	22	19	2		•••	52	56	91	289	2	1164
CLASS V—ORDER I— ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusions 2. Wounds { a Gunshot b Cuts} 3. Burns and Scalds 3a. Sunstroke 4. Poison 4a. Bite of Snake or Insect 5. Drowning 6. Suffocation 7. Otherwise	3 4 3	2 10 4 18	2 2I 2	1 4 I I I I	2 7 3 1 1	11 3 1 9 4 2 1 18	13 1 2 1 1 13	16 2 3 2 1 1 9	17 1 3 3 10	15 3 3 2 4 8 	15 1 1 14	19 1 2 1 3 10	26 I I I 9 4	19 1 3 1 1 11 	19 1 1 1 8 1	18 1 1 2 3 1	9 1 4 2 3 1	1 1 1	2 I 2	8 4	6 5	236 13 8 70 31 24 4 164 21 2
	22	34	40	8	14	50	31	34	35	36	31	36	42	37	32	26	20	16	5	12	12	573
ORDER 2.—VIOLENT DEATHS IN BATTLE. I. Gunshot Wounds 2. Sword and Bayonet Wounds. 3. Otherwise																						
ORDER 3.—HOMICIDE. Murder and Manslaughter	3				I	I	2		I			5	I		ı						2	17
ORDER 4.—SUICIDE. 1. Wounds { Gunshot Cut, Stab, &c. 2. Poison 3. Drowning 4. Hanging 5. Otherwise									 I 2	 I I 	3 I	 I I 2 	 2 1 	1 2 2 5	2 2 4	 I I 	 I 	 I 		 I 	 	1 9 8 2 10 I
ORDER 5-EXECUTION.								-														
Violent Deaths (not classed)																						
Total Class V	- 25	34	40	8	15	51	33	34	38	39	35	46	46	42	37	28	21	17	6	13	15	623
Causes not specified or ill- defined.					I	5	4		2		I	3	6	2		I	2	3			7	37
GRAND TOTALS	1547	523	305	213	142	431	212	174	218	224	256	303	302	282	246	235	313	309	276	458	48	7017

H.

SUMMARY of DEATHS of BOTH SEXES registered in the Colony of New South Wales, during each Month of the Year 1876, classified under the heads of the several Causes of Death.

	Janu	ary.	Febr	uary.	Marc	eh.	Apr	il.	Ma	ay.	Jur	ne.	- Jul	ly.	Aug	ust.	Septer	nber.	Octo	ber.	Nove	mber.	Decen	nber.	То	tal.	al.	e on hs of
CAUSES OF DEATH.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Males.	Females.	Grand Total.	Percentage on total Deaths of he Year.
I.—Zymotic Diseases. 1. Miasmatic Diseases—(Scarlatina, Diphtheria, Dysentery, Fevers, &c.) 2. Enthetic Diseases—(Syphilis, Gonorrhœa, &c.) 3. Dietic Diseases—(Privation, Scurvy, Delirium Tremens, &c.) 4. Parasitic Diseases—(Thrush, Worms, &c.) Total	18	 8 2	13 4	5 2	21	4 2	15	4 8 6	2 14 2	6 2	4 15 3	2 4 I	13	9 1	13 2	2 4 I	16	3 8 1	4	4	1 5 2	 IO I	20	5 3	33 184 23	16 75 23	49 259	26°36 0°44 2°31 0°42
II.—Constitutional Diseases. 1. Diathetic Diseases—(Gout, Dropsy, Cancer, &c.) 2. Tubercular Diseases—(Scrofula, Phthisis, Hydrocephalus, &c.) Total	10 52	20	23 41	13		15	2 I 42	19	21	20	26 54	4 3.1	20	16 12	16 44	15	19	13	13	7 24	15	13	21	8 29	238 495	163	401 876	3.28 3.83
III.—Local Diseases. 1. Diseases of the Nervous System — (Apoplexy, Paralysis, Insanity, Cephalitis) 2. Diseases of the Organs of Circulation—(Pericarditis, Aneurism, Heart Disease)	91	56			61						64				50 38		-											11.55
 Diseases of the Respiratory System—(Bronchitis, Pneumonia, Asthma, Pleurisy, &c.) Diseases of the Digestive Organs—(Gastritis, Enteritis, Peritonitis, Hernia, &c.) Diseases of the Urinary Organs—(Nephritis, Ischuria, Diabetes, &c.) Diseases of the Organs of Generation—(Ovarian Dropsy, Uterus 	43		30	29		22	34	35 37 7	30	32		27	28	21	31	22		25	60 31 7	38 25 8				26	713 388 140	324	712	10.09 6.36 1.80
Disease, &c.) 7. Diseases of the Joints—(Arthritis, Ostitis, Periostitis, &c.) 8. Diseases of Integumentary System—(Phlegmon, Ulcer, Skin Disease, &c.) Total			2			I	3		I	3	2		I	2	I			4	I			I	 180		I4 I2	6	20	0°24 0°18 0°20
IV.—Developmental Diseases. 1. Diseases of Children—(Cyanosis, Teething, &c.) 2. Diseases of Adults—(Paramenia, Childbirth, &c.) 3. Diseases of Old People—(Old Age, &c.) 4. Diseases of Nutrition—(Atrophy, Debility) Total	23 34		24 25	13 26	26 32 37 95	10 17 26	37 35	15 17 31	43	14 17 14	4I 23	17 - 16 19	50 19	18 17 23	44	12 12 14	39	13 17 9	 38 24	14 16 11	30	8 12 28	29	5 8 31	430	141	141 610 557	4'23 1'26 5'45 4'98
V.—Violence. 1. Accident or Negligence—(Fractures, Contusions, Burns, Drowning, Suffocation, Wounds, &c.) 2. Homicide—(Murder and Manslaughter) 3. Suicide—(Poison, Drowning, Hanging, &c.) 4. Execution—(Hanging) Total	5° 4 7		3 4			2	3	2	3			 I	3 2		3		46 5 	1 2 	3	2	5 3				38	7	27 47 3	
Unspecified	4	4	3	3	8	I	5	2	9	9	6	3	5	4	2	2	4	2	6	6	3	I	6	6	58	43	101	0,00

Table showing the Deaths which have occurred from the Causes stated, in each of the several Wards of the City of Sydney, from 1st January to 31st December, 1876, distinguishing those under Five years of age.

	G	ipps	S.	В	ourk	te.	Br	isba	ine.	Mac	equa	rie	(look.		Fi	itzro	у.	Ph	illip.	D	enis	on.	To	otal 1	Deatl	hs fro	om e	ach o	ause	in S	ydne	·J.
CAUSES OF DEATH.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Total.	Under.	Over.	Total.	1876.	1875.	1874.	1873.	1872.	1871.	1870.	1869.	1868.	1867.
CLASS I—ORDER I.																	,			1	1	ĺ.				,							
1. Small-pox 1a. Chicken-pox 2. Measles																																	
3. Scarlatina	9	3	12	4	4	8	7	4	II	6	8	14	42	12	54	30	19	49	15	9 2	4 2	3 11	34	206	102	37	5	54	2	3	19	.,	27
3a. Diphtheria				 I	Ι	I	2	 I		3		3	4				2				2		I	17	-		16				14	34	36
5. Croup 6. Whooping-cough	6	I	7	I		I	I		I	4		4	9		9	5	I	6	2.		2	4	4	34	27	16	18	18	5	8	-		28
7. Typhoid (and Infantile Fever.)		6	6			17			8	2	Ι,	3	3	14	17	2	9	II					9		75	39			63		46 72		34 71
8. Erysipelas		I	I			2	2				a.		2	6	6		·			2	2				18		9		8	2 7 I	8	2 12	
II. Influenza	I		I		9	9	I		I				I		I	I		I			· ··				6	2	I	3	3	2	6	4	3
12. Dysentery						20	14		 17	8	1	I	I 23	2	I 25	I I2						2 2	4	15		150		23			73		5 ²
14. Cholera	2	I	- 3							3	I	4	3		3	I		I	I	3	4 :	2 I	3	18	20	17	13	12	7	7	6	7	7
15. Ague 16. Remittent Fever																					4	2		2 4	6		3	 I			2		7
17. Rheumatism						3	1	I	I		I	I		I	I		2	2		2	2			II		9			10		6	6	0
18. Pyæmia 19. Parotitis																	2							5	6		2	3	3				
	27	19	46	10	69	79	28	20	48	26	15	41	88	39	127	57	40	97	35 2	7 6	2 46	5 26	72	572	748	440	279	345	256	237	307	383	906
ORDER 2.					,		0		0		1												-			6		7.0		4	_	-	_
 Syphilis		3	3		4								2					···			2			11	/		13		4	4		5	5
Urethra.		3	3		7	7	2		2				2		2				2 .		2	 I	I	17	I 2	9	14	15	5	5	6	12	5
Oppur																													1		9		
ORDER 3. 1. Privation	2	···	2	 I			2		2							7		7		I . I	I		2	I 38		I 22	4	31	2 I2	4 7	1 8	8	4
3. Purpura and Scurvy 4. Alcoholism:—					· I	I											I	I						2	2	4	I	2	2	2	I	3	I
a Del. Tremens b Intemperance		2	2		2 2 1				5		I	I		I 2	1 2		2 I	2 I		I .	· · · · · · · · · · · · · · · · · · ·			7 35	17 20	7 8	16	5	5	5	12	5	3 2
	2	2	4	I	24	25	2	5	7	5	2	- 7	9	3	12	7	4	II	10	2 I	2 2	3	5	83	68	42	57	45	22	24	30	27	17
ORDER 4.																									7	1	6	5	. 6	7	I	I	3
2. Worms, &c.													I		I		114							I	2	3	2	I	I			2	4
	-	-			2	2						7	I		I					I	ı			4	9	7	8	6	7	7	3	3	7
Total, Class I	_ 29	24	53	II	102	113	32	<u>-</u>	57	31	17	48	100	42	142	64	44	108	47 3	0 7	7 49	29	78	676	837	498	358	411	290	273	346	425	935
CLASS II-ORDER 1.								-				_								-				-		-		4	т	2	4	т.	I
1. Gout	3	4	7		13	13	I	5	6	I		I	2	6	8	I	4	5	I	2	3 2	5	7	50	3	9	21	16	9	22	9	17	19
3. Cancer																		8			3		5	48						28	19	20	19
3b. Polypus																																	
4. Noma																									4	4	2	7	1	2	Ι	4	4
,	4	7	II	-	34	34	I	10	II	2	7	9	2	II	13		13	14	I	5	6 :	2 10	12	IIO	71	65	85	65	38	56	33	49	43
ORDER 2.																		1							0				0	0	-		
 Scrofula Tabes Mesenterica 		I	I		I	5	I		I	4		4	I		I	0	. 2 I	2 IO			51 8	8	8	7 60	57	12	16	38	36	33	23	20	25
3. Phthisis		13	13		49	49	3	9	12		17	17	I	31	32	2	30	40	2	3 2	3 :	2 14	10	202	201	105	153	155	139	122	101	115	142
3a. Hæmoptysis	2		2	I		I	I		I	4		4	3		3	3		3	4.		4	Ι	I				7				14	4	12
5. Abscess		I	I																				I	I				I					•••
	_	_		_			_	-		-													26					-					
Total, Class II	II	22	33	6	86	92	10	20	30	10	24	34	26	42	68	15	54	69	10 2	8 3	8 1.	4 24	38	402	356	307	287	283	236	238	181	202	233
CLASS III—ORDER I. I. Cephalitis		4	4		17	17		2	2		5	5	13	6	16 6 7		5	5 5 5		4	4	. 2	5 2 4	45	67	56	55	40	44	42	40	43 18	16
3. Paralysis 4. Insanity					-												2	2					1			2	I				I	I	

I-continued.

		Jipp	os.	В	ourk	e.	Br	isba	ne.	Ma	equa	arie		Cook		F	itzroj	y.	Ph	nillij	p.	Dei	niso	n.	T	otal]	Deat	hs fro	om ea	ach c	ause	in Sy	ydne	у.
CAUSES OF DEATH.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	1876.	1875.	1874.	1873.	1872.	1871.	1870.	1869.	1868,	1867.
CLASS III—ORDER 1— continued. 5. Chorea 6. Epilepsy 7. Convulsions 8. Brain Disease, &c	13	2 2	13	I	10	11	10 	2 I	2 10 1	9	I 	9	29	5	1 29 7	14	 8	14 9	ī 5 	I 2 4	1 17 4		2	19	36 —	9 106 41	51	105 45	16 127 35	36	87 21	4	7 91 24	26
Order 2. 1. Pericarditis		8 -	1	3		I		12	12		6	6		16	16 ——			20	1	 I I	11		10	10	7	9 107 —-	7 91 —		8 92	93	89	7 13 84 ———————————————————————————————————	71	
Order 3. 1. Laryngitis 2. Bronchitis 3. Pleurisy 4. Pneumonia	6	7 3	13 3	3 I 3 ···	25 2 25 2	25 26 	3	2 2 3 1	12 2 6 1 1	I I I	5	 I	7 4 	8 1 2 3 1 1	7	7 4 2	3 3 10 2 3	3 14 4 	7 3 1	5 1 3 1	12 1 6 2 2	4 1	7 3 3	16 3 7 1	108 14 82 19 3 11	15 106 13 11	104 9 }94 11 8	80 4 73 9 15	80 6	4 83 1 10	6 64 3 21	1 69 11 60	16 55 6 17	7 44 5
Order 4. 1. Gastritis	· 3	1 3 1 2	3 3 1 2 2	4 I	5 8 3	5 8 	2 2	 I	2 2 I 	 I	I I	 2 I	4 6	 3 2 I	4 9	4 4	I	5 4	2 . 2	 I I 	24 2 2 1 1 	4 2		4 2		35	14 36 13 4 2 3	 5		10		13 25 16 1 2 3	4 27 9 1 3 	174 13 25 7 2 4 2 5
9. Stricture of Intestine 10. Fistula	s . I	3	3	i	 4 7 2 1	4 7 2 1	I	 I 6 I 4	 6 1 4	[1 2 5	 1 2 5 	2 I 	4 6 	3	 I	4	4 9		3 2	3 2	6	5	 1 5 1	1 15 31 7 32	2 7 23 5 51 	2 7 5 41 1	19 9 8 38	 15 15 5 26	I		2 11 10 1 36 		1 10 6 2 38
ORDER 5. 1. Nephritis 2. Ischuria 3. Nephria 4. Diabetes 5. Stone 6. Cystitis 7. Kidney Disease, &c.		5	5	5	15	15		2 I	 2 I		2	3		7 1 2 2	7 1		4	 4	2	3	5	I I 2	I 2 I 4	2 3 I		4 3 8 12	7 2 37 2 2 3 18		1 3 15	4 1 20 3 1 3 9	 4 3 1 2 8	2 2 10 2 2 9	 2 8 1 4 6	 1 2 1 2 6
ORDER 6. 1. Ovarian Dropsy 2. Uterus Disease, &c									·			2					•			I	1				1 6 7	 5 — 5	55	2 3 5	32	3 2 5	I 2 3	4	1 9	I 2 3
ORDER 7. 1. Arthritis 1a Ostitis ,Periostitis, &c 2. Joint Disease, &c					 I	 I				 I						100000000000000000000000000000000000000									3 2	2 I —		 1 3 	 	I I		3	2	2
ORDER 8. 1. Phlegmon 2. Ulcer 3. Skin Diseases, &c		I			I I 																				3	1 2	2 2	 I 2	5 3	 5 2	4 2	1 1 3	4	 I
Total Class III	32	63	95	5 5	186	191	35	48	83	21	34	55	81	90	171	43	90	133	38	54	92	44	48	92	912	982	877	787	768	732	530	647	595	512

I—continued.

	1	Fipp	S.	E	Bourk	e.	Bri	isba	ne.	Mad	equa	1		Cook		F.	itzroj	7.	P	hilli	p.	De	niso	n.	T	otal :	Deat	hs fr	om e	ach (cause	in S	ydne	y.
CAUSES OF DEATH.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	Under.	Over.	Total.	1876.	1875.	1874.	1873.	1872.	1871.	1870.	1869.	1868.	1867.
CLASS IV.—ORDER 1. 1. Premature Birth 2. Cyanosis 3. Spina Bifida 4. Other Malformations. 5. Teething	5 1 1 2		5 1 1 2	3 1		3 1	2 I		2 I 4	7		7 1 3	23 I I 		23 I I II	3 2 6		3 2 6	10 1 2 5		10 1 2 5	7 I 4		7 1 4	60 5 3 5 35	67 4 5 41	28 2 2 6 49	37 7 32	37 3 3 53	40 3 2 9 47	3 ² 4 10 37	32 7	33 4 3 39	23 I 3 56
ORDER 2. 1. Paramenia		3 4	3 4					 I I	 I		 I	I		5 5	5 5		I I 2	I I 2		5 5	5 5				2 17 19	30	4	3		3	4 4	3	I	I I 2
ORDER 4. 1. Atrophy and Debility Total Class IV CLASS V.—ORDER 1.—	. 18			-			-																											
ACCIDENT OR NEGLIGENCE. 1. Fractures & Contusion 2. Wounds { a Gunshot.} b Cuts 3. Burns and Scalds 3a.Sunstroke 4. Poison 4a.Bite of Snake or Insec 5. Drowning 6. Suffocation 7. Otherwise—Shock, &ce	s t	I 2	 I 2	2	 9 1 6 1	II II I 6	 I 	 I 3	2 3 I				 2 I	2 2	2 4 	2		 2 I	I	I I 	 I 2 				 4 22 I 12 6	 6 10 1 4 1	} 4 9 3 i 16 2	5 19 4 1 21	3 8 2 I	2 8 	3 12 1 3 16 10	2 8 1 1 14 5	 II I	6
ORDER 2.— VIOLENT DEATHS IN BATTLE 1. Gunshot Wounds 2. Sword and Bayonet Wounds. 3. Otherwise																																		
ORDER 3.—HOMICIDE. 1. Murder 2. Manslaughter ORDER 4.—SUICIDE.					3	 3 4	I		I I			:					I				2		I	 I		3 2	8 4	4 2	5 1	4 5	I	4	 I	 I
1. Wounds {Gunshot 2. Poison 3. Drowning 4. Hanging 5. Otherwise					2 I	2 I 					I 	 I		 I									I	I	4 2 1	5 4 1 3 1	3 	4 1 1	3 1 2	2	4 2	2 2 3 1 1 1		1 2
Violent Deaths (not classed) — Found Drowned, &c.		-			-	5		5	5					1	1		1	I	I		ľ		1	I	14	8								
Total Class V Causes not specified or ill defined.	-	-		-	12	12		4	4		3	3	1	9	10		12	12	2	6	8		6	6	55	19	15	18	20	18	18	25	60	54
	91	125	216	32	455	487	93	113	206	87	88	75	289	207	496	152	219	371	139	140	279	141	120	201	2491	2073	2077	1800	1810	1583	1492	1511	1599	2151